

OCT 26 1953

MEDICAL LIBRARY
HISTORICAL BUILDING
DES MOINES, IOWA

Arizona Medicine

Journal of

ARIZONA MEDICAL ASSOCIATION

**SOUTHWESTERN
MEDICAL
CONVENTION**


October 29, 30, 31, 1953



**PIONEER
HOTEL
TUCSON, ARIZONA**

ILOTYCIN, PEDIATRIC
(ERYTHROMYCIN, LILLY), ETHYL CARBONATE, CRYSTALLINE
a most effective antibiotic
for the common bacterial
infections of childhood
AVAILABLE IN 60-cc. BOTTLES

Lilly



dependable decongestion

in

colds,

sinusitis, allergic rhinitis

Through its prompt and prolonged decongestive action Neo-Synephrine not only restores nasal patency during all stages of the common cold, sinusitis or allergic rhinitis, but also helps to reestablish and protect the physiologic defense mechanisms of the nasal cavity: and encourages proper sinus drainage and aeration.

Neo-Synephrine's powerful vasoconstrictive action is exerted with virtually no sting, congestive rebound, or systemic side effects and is undiminished after repeated use.

NEO-SYNEPHRINE[®]

Hydrochloride

Brand of phenylephrine

Rapid and Sustained Relief

¼% solution (plain and aromatic),
1 oz. bottles

½ and 1% solutions (when stronger
vasoconstrictive action is needed), 1 oz.
bottles

¼% water soluble jelly, ½ oz. tubes
Neo-Synephrine, trademark reg. U.S. & Canada.

Winthrop Stearns, INC.
NEW YORK 18, N. Y. WINDSOR, ONT.

ACCIDENT**HOSPITAL
INSURANCE****SICKNESS****For Physicians, Surgeons, Dentists Exclusively**

\$5,000 accidental death Quarterly **\$8.00**
\$25 weekly indemnity, accident and sickness

\$15,000 accidental death Quarterly **\$24.00**
\$75 weekly indemnity, accident and sickness

\$10,000 accidental death Quarterly **\$16.00**
\$50 weekly indemnity, accident and sickness

\$20,000 accidental death Quarterly **\$32.00**
\$100 weekly indemnity, accident and sickness

COST HAS NEVER EXCEEDED AMOUNTS SHOWN

ALSO HOSPITAL INSURANCE

	Single	Double	Triple	Quadruple
60 days in Hospital	5.00 per day	10.00 per day	15.00 per day	20.00 per day
30 days of Nurse at Home	5.00 per day	10.00 per day	15.00 per day	20.00 per day
Laboratory Fees in Hospital	5.00	10.00	15.00	20.00
Operating Room in Hospital	10.00	20.00	30.00	40.00
Anesthetic in Hospital	10.00	20.00	30.00	40.00
X-Ray in Hospital	10.00	20.00	30.00	40.00
Medicines in Hospital	10.00	20.00	30.00	40.00
Ambulance to or from Hospital	10.00	20.00	30.00	40.00
	CO-INSURANCE (Quarterly)			
Adult	2.50	5.00	7.50	10.00
Child to age 10	1.50	3.00	4.50	6.00
Child over age 19	2.50	5.00	7.50	10.00

\$4,000,000.00
INVESTED ASSETS

PHYSICIANS CASUALTY ASSOCIATION
PHYSICIANS HEALTH ASSOCIATION

\$19,500,000.00
PAID FOR CLAIMS

51 years under the same management

400 First National Bank Building

Omaha 2, Nebraska


\$200,000.00 deposited with State of Nebraska for protection of our members

SOUTHWESTERN SURGICAL SUPPLY CO.

YOUR COMPLETE SOURCE IN THE SOUTHWEST
 FOR ALL ETHICAL MEDICAL EQUIPMENT AND
 SUPPLIES.

**PHOENIX****TUCSON****ALBUQUERQUE****EL PASO**

Cervical *and* Thoracic Regions *of the* Spinal Cord



This is one of a series of paintings for Lederle by Paul Peck, illustrating the anatomy of various organs and tissues of the body which are frequently attacked by infection, where aureomycin may prove useful.



*A great many varieties
of pathogens have been proved
susceptible to*

Aureomycin

HYDROCHLORIDE CRYSTALLINE

*The rapid diffusion of
Aureomycin into the
cerebrospinal fluid makes
it a drug of choice for
the treatment of meningitis.*

LEDERLE LABORATORIES DIVISION AMERICAN Cyanamid COMPANY
30 Rockefeller Plaza, New York 20, N. Y.

ARIZONA MEDICINE

Journal of ARIZONA MEDICAL ASSOCIATION

VOL. 10, NO. 10



OCTOBER, 1953

TABLE OF CONTENTS

OFFICERS

ARIZONA MEDICAL DIRECTORY	VI
WOMAN'S AUXILIARY DIRECTORY	VI
COUNTY MEDICAL SOCIETY OFFICERS DIRECTORY	VIII

ORIGINAL ARTICLES

INDICATIONS FOR IRRADIATION OF BREAST LESIONS	353
U. V. Portmann, M.D., Tucson, Arizona	
THE ROLE OF THE NON-SPECIALIST IN THE CARE OF ALLERGIC DISEASE....	354
Samuel M. Feinberg, M.D., Chicago, Illinois	
WHY REPORT COMMUNICABLE DISEASES?	358
Drs. Clarence G. Salsbury and James B. Swayne, Phoenix, Arizona	

MEDICAL PROBLEMS

PHOENIX CLINICAL CLUB	363
Drs. Louis B. Baldwin and A. J. Fillmore	

PRESIDENT'S PAGE

SPEARS CHIROPRACTIC HOSPITAL IN DENVER	372
--	-----

EDITORIAL

HOSPITAL BEDS	373
---------------------	-----

ARIZONA PHARMACEUTICAL PAGE

THE SECOND CENTURY 1852-1953	374
------------------------------------	-----

TOPICS OF CURRENT MEDICAL INTEREST

RX., DX., AND DRS.	376
Guillermo Osler, M.D.	
DIFFERENTIAL DIAGNOSIS IN THE RHEUMATIC DISEASES	371
INTERNATIONAL ACADEMY OF PROCTOLOGY	371
MEETING NOTICE	371
CHIROFODY, A HAND MAIDEN OF MEDICINE	381

WOMAN'S AUXILIARY

THE PLEDGE OF LOYALTY TO THE WOMAN'S AUXILIARY	380
Mrs. David W. Thomas	
LEGISLATION REPORT	380
Mrs. Charles L. von Pohle, Chandler, Arizona	

DIRECTORY

LABORATORIES	XXI
SANATORIUM DIRECTORY	XXII
DRUGGISTS DIRECTORY	XXVI
PHYSICIANS DIRECTORY	XXIX

Published monthly by the Arizona Medical Association. Business office at 424 Heard Building, Phoenix, Arizona. Subscription \$3.00 a year, single copy 25c. Entered as second class matter March 1, 1921, at Postoffice at Phoenix, Arizona, Act of March 3, 1879.

Directory

THE ARIZONA MEDICAL ASSOCIATION, INC.
Organized 1892 341 SECURITY BUILDING
234 NORTH CENTRAL AVE., PHOENIX, ARIZONA
OFFICERS AND COUNCIL

Edward M. Hayden, M.D. President
1603 N. Tucson Boulevard, Tucson, Arizona
Oscar W. Thoeny, M.D. President-Elect
1313 N. 2nd Street, Phoenix, Arizona
Harry E. Thompson, M.D. Vice-President
433 N. Tucson Boulevard, Tucson, Arizona
Dermont W. Melick, M.D. Secretary
15 East Monroe, Phoenix, Arizona
Clarence E. Yount, Jr., M.D. Treasurer
P. O. Box 1626, Prescott, Arizona
Lindsay E. Beaton, M.D. Speaker of House
1650 N. Campbell Ave., Tucson, Arizona
Jesse D. Hamer, M.D. Delegate to AMA
15 East Monroe, Phoenix, Arizona
Robert E. Hastings, M.D. Alternate-Delegate to AMA
1014 N. Country Club Road, Tucson, Arizona
R. Lee Foster, M.D. Editor-In-Chief
15 East Monroe, Phoenix, Arizona

DISTRICT COUNCILORS

David C. Jamra, M.D. Central District
926 E. McDowell Rd., Phoenix, Arizona
Kent H. Thayer, M.D. Central District
1313 N. 2nd Street, Phoenix, Arizona
Leslie B. Smith, M.D. Central District
926 East McDowell Road, Phoenix, Arizona
Herbert B. Potthoff, M.D. Northeastern District
P. O. Box 128, Holbrook, Arizona
Ernest A. Born, M.D. Northwestern District
107 North Cortez, Prescott, Arizona
Guy B. Atonna, M.D. Southeastern District
447 10th Street, Douglas, Arizona
Wilkins R. Manning, M.D. Southern District
620 N. Country Club Road, Tucson, Arizona
Royal W. Rudolph, M.D. Southern District
1627 N. Tucson Boulevard, Tucson, Arizona
Abe I. Podolsky, M.D. Southwestern District
1601 5th Avenue, Yuma, Arizona
COUNCILOR AT LARGE
Thomas H. Bate, M.D. Past President
15 East Monroe, Phoenix, Arizona

BOARDS

PROFESSIONAL: Hugh C. Thompson, M. D., Tucson (Chairman); Ernest A. Born, M. D., Prescott; Edward H. Bregman, M. D., Phoenix; Orin J. Farness, M. D., Tucson; Lewis H. Howard, M. D., Tucson; Elvie B. Jolley, M. D., Bisbee; Joseph M. Kinkade, M. D., Tucson; Milton C. F. Semoff, M. D., Tucson; George A. Williamson, M. D., Phoenix.
PUBLIC RELATIONS: Leo J. Kent, M. D., Tucson (Chairman); Howard C. Lawrence, M. D., Phoenix (Vice Chairman); Lindsay E. Beaton, M. D., Tucson; Carlos C. Craig, M. D., Phoenix; Donald F. DeMarse, M. D., Holbrook; Wilkins R. Manning, M. D., Tucson; Robert M. Matts, M. D., Yuma; Zenas B. Noon, M. D., Nogales.

STANDING COMMITTEES

GRIEVANCE: Thomas H. Bate, M. D., Phoenix (Chairman); Lindsay E. Beaton, M. D., Tucson; Preston T. Brown, M. D., Phoenix; Arthur C. Carlson, M. D., Phoenix; Robert E. Hastings, M. D., Tucson; Hilary D. Ketcherside, M. D., Phoenix; Harry T. Southworth, M. D., Prescott.

HISTORY & OBITUARIES: W. Warner Watkins, M. D., Phoenix (Historian); R. Lee Foster, M. D., Phoenix; Dermont W. Melick, M. D., Phoenix; Hal W. Rice, M. D., Tucson.

INDUSTRIAL RELATIONS: William B. McGrath, M. D., Phoenix (Chairman); Carl H. Gans, M. D., Morenci; Ronald S. Haines, M. D., Phoenix; Zenas B. Noon, M. D., Nogales; John R. Schwartzmann, M. D., Tucson.

LEGISLATION: Reed D. Shupe, M. D., Phoenix (Chairman); Jesse D. Hamer, M. D., Phoenix (Vice Chairman); Carl W. Ahl, M. D., Douglas; William H. Bates, M. D., Cottonwood; Walter Brazie, M. D., Kingman; Ellis V. Browning, M. D., Springerville; Max Costin, M. D., Tucson; Donald F. Hill, M. D., Tucson; Bert E. Lambrecht, M. D., Miami; Stanton C. Love, M. D., Clifton; Wilkins R. Manning, M. D., Tucson; Robert M. Matts, M. D., Yuma; Zenas B. Noon, M. D., Nogales; Donald A. Polson, M. D., Phoenix; Wallace A. Reed, M. D., Phoenix; Leo Schnur, M. D., Grand Canyon; Leslie B. Smith, M. D., Phoenix; Noel G. Smith, M. D., Safford; Guss B. Steward, M. D., Coolidge; Kent H. Thayer, M. D., Phoenix; Harry E. Thompson, M. D., Tucson; Myron G. Wright, M. D., Winslow.

MEDICAL DEFENSE: Preston T. Brown, M. D., Phoenix (Chairman); Ernest A. Born, M. D., Prescott; Harold W. Kohl, M. D., Tucson.

MEDICAL ECONOMICS: Hilary D. Ketcherside, M. D., Phoenix (Chairman); Royal W. Rudolph, M. D., Tucson; Hugh C. Thompson, M. D., Tucson.

PUBLISHING: R. Lee Foster, M. D., Flagstaff; Donald E. Nelson, M. D., Safford; Darwin W. Neubauer, M. D., Tucson.

SCIENTIFIC ASSEMBLY: Oscar W. Thoeny, M. D., Phoenix (Chairman); Joseph Bank, M. D., Phoenix; David Engle, M. D., Tucson; Francis M. Findlay, M. D., Kingman; David C. Jamra, M. D., Phoenix; Wilkins R. Manning, M. D., Tucson; Donald E. Nelson, M. D., Safford; William A. Phillips, M. D., Yuma; Herbert B. Potthoff, M. D., Holbrook.

Woman's Auxiliary

OFFICERS OF THE AUXILIARY TO THE ARIZONA
MEDICAL ASSOCIATION - 1953 - 1954

President Mrs. George S. Enfield
335 W. Cambridge Ave., Phoenix
President-Elect Mrs. Brick P. Storts
3228 E. 5th Street, Tucson
1st Vice President (Organization) Mrs. Roy Hewitt
130 Camino Miramonte, Tucson
2nd Vice President (Program) Mrs. Charles S. Powell
698 Ninth Ave., Yuma

Treasurer Mrs. R. Lee Foster
2215 N. 11th Ave., Phoenix
Recording Secretary Mrs. Louis Hirsch
Rt. 6, Box 710, Tucson
Corresponding Secretary Mrs. William F. Schoffman
36 N. Country Club Dr., Phoenix
Director (1 year) Mrs. T. C. Harper
175 S. Third Street, Globe
Director (1 year) Mrs. William F. Schoffman
36 N. Country Club Dr., Phoenix
Director (2 year) Mrs. Frederick W. Knight
Safford

STATE COMMITTEE CHAIRMEN 1953-1954

Bulletin Mrs. Hiram Cochran
2716 E. 4th Street, Tucson
Chaplin Mrs. James R. Moore
395 W. Granada, Phoenix
Civil Defense Mrs. Frank A. Shallenberger
345 S. Eastbourne Dr., Tucson
Convention Mrs. A. E. Cruthirds
60 E. Country Club Drive, Phoenix
Finance Mrs. Paul S. Causey
38 W. Camelback Road, Phoenix
Historian Mrs. Benjamin Herzberg
125 W. Rose Lane, Phoenix
Legislation Mrs. C. L. Von Pohle
274 N. Arizona Ave., Chandler
Medical Education Fd. Mrs. J. B. Tucker
Box 1042, Casa Grande
Nurse Recruitment Mrs. Max Costin
2648 E. 4th Street, Tucson
Parliamentarian Mrs. Royal W. Rudolph
3935 Miner, Tucson
Publicity Mrs. Ashton B. Taylor
46 E. Marlette Dr., Phoenix
Public Relations Mrs. John K. Bennett
2325 E. Waverly St., Tucson
Revisions Mrs. Jesse D. Hamer
1819 N. 11th Ave., Phoenix
Student Nurse Loan Fund Mrs. Donald A. Polson
2163 W. Earl Dr., Phoenix
Today's Health Mrs. James C. Soderstrom
Box 82, Whipple
Nominating Com. Chairman Mrs. Royal W. Rudolph
3935 Miner, Tucson
Mental Health Mrs. Delbert Secrist
35 Calle Clara Vista, Tucson

COUNTY PRESIDENTS AND OFFICERS 1953-1954


GILA COUNTY
President Mrs. Ira E. Harris
Box 1296, Miami
Vice President Mrs. Charles T. Collopy
Box 623, Miami
Secretary-Treasurer Mrs. William E. Bishop
605 S. 3rd St., Globe
MARICOPA COUNTY
President Mrs. Paul B. Jarrett
501 E. Pasadena, Phoenix
President-Elect Mrs. Robert H. Cummings
5830 Arcadia Lane, Phoenix
1st Vice President Mrs. C. Selby Mills
5821 N. 11th, Phoenix
2nd Vice President Mrs. George A. Williamson
537 W. Rose Lane, Phoenix
Rec. Secretary Mrs. Robert A. Price
335 W. Windsor Ave., Phoenix
Cor. Secretary Mrs. Samuel H. Hale
Rt. 1, Box 382, Scottsdale
Treasurer Mrs. Dwight H. Porter
35 N. 12th, Phoenix








GRAHAM COUNTY
President Mrs. Noel Grey Smith
Safford
Vice-Pres. Mrs. Robert Keller
Safford
Sec.-Treas. Mrs. Donald E. Nelson
Safford

PIMA COUNTY
President Mrs. Hugh C. Thompson
135 Palomar Dr., Tucson
President-Elect Mrs. Joseph M. Kinkade
335 S. Country Club Rd., Tucson
1st Vice President Mrs. Seymour I. Shapiro
2410 E. Elm St., Tucson
2nd Vice President Mrs. George A. Shetter
2733 E. Mable St., Tucson
Rec. Secretary Mrs. John K. Bennett
2325 E. Waverly St., Tucson
Cor. Secretary Mrs. D. E. Engle
4012 E. Cooper, Tucson
Treasurer Mrs. Richard K. Hausmann
2639 E. 8th St., Tucson

PINAL COUNTY
President Mrs. Howard Holmes
Eloy
Vice President Mrs. James T. O'Neil
Casa Grande
Sec.-Treasurer Mrs. Robert Z. Collings
Box 475, Casa Grande
YAVAPAI COUNTY
President Mrs. Melvin W. Phillips
829 Flora Ave., Prescott
Vice President Mrs. Earnest Geever
Whipple
Sec.-Treasurer Mrs. William A. Marlowe
Whipple

YUMA COUNTY
President Mrs. Phillip G. Corliss
Somerton, Box 68
Vice President Mrs. John J. Kloby
Ave. C, Yuma
Sec.-Treasurer Mrs. Robert M. Matts
Yuma

cortisone +  = Cortogen

The symbol  denotes leadership in steroid hormone  research and manufacture -- including Cortogen in the most useful clinical forms. For systemic therapy, use Cortogen tablets  or injection . For eye  disorders use Cortogen ophthalmic suspension -- sterile . The Schering seal  appears on each tablet and package -- your assurance of quality.

CORTOGEN[®]

Acetate (cortisone acetate, Schering) Tablets, 5 mg. and 25 mg.;
Injection, 25 mg. per cc., 10 cc. multiple-dose vials;
Ophthalmic Suspension—Sterile, 0.5% and 2.5%, 5 cc. dropper bottles.

Schering

CORPORATION • BLOOMFIELD, NEW JERSEY
In Canada: Schering Corporation, Ltd., Montreal.

CORTOGEN



THE ARIZONA MEDICAL ASSOCIATION, INC.
COUNTY MEDICAL SOCIETY OFFICERS FOR 1953

APACHE COUNTY		MOHAVE COUNTY	
President	Ellis V. Browning, M.D.	President	Walter Brazie, M.D.
Secretary	Springerville	Secretary	Masonic Building, Kingman
	Springerville		Francis M. Findlay, M.D.
	Springerville		Masonic Building, Kingman
COCHISE COUNTY		NAVAJO COUNTY	
President	Carl W. Ahl, M.D.	President	Myron G. Wright, M.D.
Secretary	610 Ninth St., Douglas	Secretary	122 W. 3rd St., Winslow
	George N. Watson, M.D.		Herbert B. Potthoff, M.D.
	620 Tenth St., Douglas		Box 128, Holbrook
COCONINO COUNTY		PIMA COUNTY	
President	Leo Schnur, M.D.	President	Wilkins R. Manning, M.D.
Secretary	Box 115, Grand Canyon	Secretary	620 N. Co. Club Rd., Tucson
	Charles W. Sechrist, M.D.		William B. Steen, M.D.
	1100 N. Beaver, Flagstaff		110 S. Scott, Tucson
GILA COUNTY		PINAL COUNTY	
President	Bert E. Lambrecht, M.D.	President	Guss B. Steward, M.D.
Secretary	Box 1837, Miami	Secretary	321 W. Central, Coolidge
	Cyril M. Cron, M.D.		Chalmers D. Johnson, M.D.
	Box 1207, Miami		321 W. Central, Coolidge
GRAHAM COUNTY		SANTA CRUZ COUNTY	
President	Noel G. Smith, M.D.	President	Zenas B. Noon, M.D.
Secretary	616 12th St., Safford	Secretary	Gebler Building, Nogales
	Morris T. Sandeno, M.D.		Charles S. Smith, M.D.
	803 7th St., Safford		Gebler Building, Nogales
GREENLEE COUNTY		YAVAPAI COUNTY	
President	Stanton C. Lovre, M.D.	President	William H. Bates, M.D.
Secretary	Clifton	Secretary	Marcus Lawrence Hospital, Cottonwood
	Bernard L. Hicks, M.D.		Clarence E. Yount, Jr., M.D.
	Morenci Hospital, Morenci		Box 1626, Prescott
MARICOPA COUNTY		YUMA COUNTY	
President	Kent H. Thayer, M.D.	President	Robert M. Matts, M.D.
Secretary	1313 N. 2nd St., Phoenix	Secretary	1601 Fifth Ave., Yuma
	Wallace A. Reed, M.D.		Robert E. Rider, M.D.
	926 E. McDowell Rd., Phoenix		167 E. 3rd St., Yuma



We offer the benefit of our years of
experience in
real estate transactions

CLEVENGER REALTY

Jack Clevenger, Realtor

W. Thomas at Central Avenue

Phone AM 6-2494

Phoenix

Arizona

"Stands Ready To Serve"

We Solicit Your Business

We base this solicitation on our long years of experience in handling the investment accounts of several thousand men and women in Phoenix and the State of Arizona.

Our personnel is well qualified to render you intelligent service.

Our facilities are complete and enable us to offer you excellent service in all departments of investing — government and tax free municipal bonds; corporation stocks and bonds; local securities; listed or "over-the-counter" issues; new underwritings; investment trusts.

You will also find our ground floor location in the heart of Phoenix a great convenience with free parking at Phoenix Title & Trust parking lot, Second Avenue at Monroe.

We believe you will like doing business with us.

Phone AL 8-6646


REFSNES, ELY, BECK & CO.

Members New York Stock Exchange
112 West Adams Street — Phoenix, Arizona

HydroCortone®

(HYDROCORTISONE, MERCK)

*offers definite
therapeutic
advantages*



Primary Sites of Pathology and Indications

1. **EYE**—Inflammatory eye disease. 2. **NOSE**—Intractable hay fever. 3. **LARYNX**—Laryngeal edema (allergic). 4. **BRONCH**—Intractable bronchial asthma. 5. **LUNG**—Sarcoidosis. 6. **HEART**—Acute rheumatic fever with carditis. 7. **BONES, JOINTS, AND BURSÆ**—Osteoarthritis; Rheumatoid arthritis; Rheumatoid spondylitis; Acute gouty arthritis; Still's disease; Psoriatic arthritis; Bursitis. 8. **SKIN AND CONNECTIVE TISSUE**—Pemphigus; Disseminated lupus erythematosus; Scleroderma (early); Dermatomyositis; Atopic dermatitis; Exfoliative dermatitis; Dermatitis venenata (e.g., poison ivy); Dermatitis medicamentosa. 9. **ADRENAL GLAND**—Congenital adrenal hyperplasia; Addison's disease; Following adrenalectomy for hypertension, Cushing's syndrome, and neoplastic diseases. 10. **BLOOD, BONE, AND MARROW**—Allergic purpura; Acute leukemia* (lymphocytic or granulocytic); Chronic lymphatic leukemia.* 11. **LYMPH NODES**—Lymphosarcoma; * Hodgkin's disease.* 12. **ARTERIES AND CONNECTIVE TISSUE**—Periarthritis nodosa (early). 13. **KIDNEY**—Nephrotic syndrome, without uremia (to induce withdrawal diuresis). 14. **VARIOUS TISSUES**—Angioneurotic edema; Serum sickness; Sarcoidosis; Drug sensitization; Waterhouse-Friderichsen syndrome.

*Transient beneficial effects.

1
HYDROCORTONE is a natural and principal anti-inflammatory adrenocortical steroid.

2
HYDROCORTONE Tablets produce the same therapeutic results as cortisone, and in smaller dosage.

3
HYDROCORTONE Tablets generally may be administered in a dosage two-thirds that of cortisone.

4
HYDROCORTONE Tablets recently were drastically reduced in price. Cost of therapy now is substantially the same as with cortisone.

Literature on request

HYDROCORTONE is the registered trade-mark of Merck & Co., Inc. for its brand of hydrocortisone.



MERCK & CO., INC.
Manufacturing Chemists
RAHWAY, NEW JERSEY

© Merck & Co., Inc.

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

A Family SAVINGS Institution



Mother, Dad, Sister and Brother . . . all invited to open and add regularly to an INSURED First Federal Savings account.

Everyone in the family will enjoy the cheerful service, 3% earnings and INSURED safety.

First Federal Savings

ACCOUNTS INSURED

Earnings

3%

per annum

PHOENIX: 30 West Adams Street
2933 N. Central • 4201 S. Central
YUMA: Orange at Fourth Avenue

MR

MEDICAL RENTS

- HOSPITAL
BEDS
FRACTURE
BEDS



- WHEEL
CHAIRS
HOSPITAL
& PORTABLE



- INVALID
WALKERS
ADULT & CHILD SIZE



OXYGEN THERAPY SERVICE
Phone ALpine 4-9227
Phoenix, Arizona

Rainbow Water

☆

A constantly reliable bottled water . . .
Pure . . . Fresh . . . Naturally Soft
Untreated . . . Sterilized Equipment
Delivered. Also Distilled Water.

☆

PHONE 2-4645

☆

RAINBOW WATER CO.

332 East Seventh

Tucson

WHEN AN ORTHOPEDIC MATTRESS IS INDICATED

Restful, healthful body adjustment is supplied by the Spring Air Back Supporter Mattress, with its high density construction of lightly compressed coils of extra large diameter. Made of conventional, time-proven materials, to a new design which provides positive back support without interfering with circulation. See it at your favorite furniture store . . . recommend it with confidence.



Manufactured in Phoenix by
SOUTHWEST MATTRESS CO.
1710 EAST WASHINGTON ST.
PHOENIX, ARIZONA



an agent of choice in urinary tract infections

- promptly effective against a broad-spectrum of urinary pathogens
- high concentration in active form in urinary tract
- well tolerated, even upon prolonged administration

Terramycin
is acclaimed
by urologists everywhere
for unsurpassed action in
*chronic urinary tract
infections*
*acute urinary tract
infections*
urinary tract surgery

Pfizer

PFIZER LABORATORIES, INC.



RAYTHEON

Microtherm

wherever diathermy is indicated

RAYTHEON Radar *Microtherm* offers you the modern microwave method of precision heat application.

MICROTHERM operates at 2450 megacycles, as contrasted with the highest television range of 920 megacycles, hence TV interference is avoided.

MICROTHERM provides penetrating energy for deep heating — dosage may be accurately timed.

MICROTHERM is safe as well as quick, easy to apply as well as clinically efficient.

Ask for a demonstration or let us mail you the latest clinical reports on Radar Microwave Diathermy.

APPROVED BY THE F. C. C.
CERTIFICATE NO. D-477
UNDERWRITERS LABORATORY



RAYTHEON
Leaders in Electronics

RAYTHEON MANUFACTURING COMPANY
Power Tube Division - Waltham 54, Mass.

TAKE THE TIME to investigate the diathermy equipment used in leading clinics, hospitals and doctors' offices — over fifteen thousand *Microtherms* now in use.




STANDARD SURGICAL SUPPLY
710 NORTH FIRST STREET PHOENIX • ARIZONA

BRANCH: 1706 E. SPEEDWAY — TUCSON, ARIZONA

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

"PREMARIN"

...*"sense of well-being"*...

In addition to relief of menopausal symptoms,
"a feeling of well-being or tonic effect" was frequently
reported by patients on "Premarin" therapy.*

"PREMARIN®" in the menopause

Estrogenic Substances (water-soluble) also known as
Conjugated Estrogens (equine). Tablets and liquid.

*Harding, F. E.: West. J. Surg. 52:31 (Jan.) 1944.

"PREMARIN"

AYERST, MCKENNA & HARRISON LIMITED • New York, N. Y. • Montreal, Canada

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

A Wise Choice Against Resistant Cocci

ERYTHROCIN*



a selective antibiotic



A DRUG OF CHOICE

against the majority of coccic infections—especially when patients are sensitive to other antibiotics or the cocci are resistant.



ADVANTAGEOUS

because the special acid-resistant coating, developed by Abbott, and Abbott's built-in disintegrator, assure rapid dispersal and absorption in the upper intestinal tract.



A DRUG OF CHOICE

against staphylococci—because of the high incidence of staphylococcic resistance to other antibiotics.



A DRUG OF CHOICE

because it is less likely to alter normal intestinal flora than other oral antibiotics, except penicillin; gastrointestinal disturbances are rare; no serious side effects reported.




Prescribe ERYTHROCIN

in pharyngitis, tonsillitis, otitis media, sinusitis, bronchitis, pneumonia, scarlet fever, erysipelas, pyoderma, certain cases of osteomyelitis, and other indicated conditions.

Abbott

* Trade Mark for

**ERYTHROMYCIN, ABBOTT
CRYSTALLINE**

RAPID ABSORPTION—MAXIMUM THERAPEUTIC EFFECT

The clinical effectiveness of different brands of mephenesin tablets depends on their rate of absorption. A mephenesin tablet that disintegrates slowly is absorbed slowly. The resulting low blood levels may never produce a maximum therapeutic effect. Results with such a tablet are usually poor.

Tolserol Tablets are a result of extensive study and are formulated to disintegrate rapidly for fast absorption, thus maintaining optimum blood levels.

Tolserol

(Squibb Mephenesin)

Complete information on the use of Tolserol in muscle spasm of rheumatic disorders, in neurologic disorders and in acute alcoholism is available from the Professional Service Department, Squibb, 745 Fifth Avenue, New York 22, N. Y.

SQUIBB

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL


Upjohn

absorbable hemostat:

Available in a large variety of sizes and forms, including:

Surgical sponges
Compressed surgical sponges
Dental packs
Gynecologic packs
Nasal packs
Prostatectomy cones
Tumor diagnosis kit

The Upjohn Company, Kalamazoo, Michigan



Gelfoam



Trademark Reg. U.S. Pat. Off. BRAND OF ABSORBABLE GELATIN SPONGE

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

Here's the most comprehensive x-ray supply catalog ever published!



No x-ray department can afford to be without General Electric's new x-ray supply catalog. Every supply and accessory item you need is covered in an easy, straight-forward manner that simplifies ordering.

And here are two unique conveniences: Prices are printed alongside every listing — there's no need to bother with a separate price list. Bound-in postpaid order cards also save time — and postage.

Ask your G-E x-ray representative for this handy reference guide to your entire x-ray supply needs.

GENERAL  ELECTRIC

Direct Factory Branch: 821 West Adams, PHOENIX

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

New Horizons in Antibiotic Therapy

BICILLIN[®]

Dibenzylethylenediamine Dipenicillin G

A NEW FORM OF PENICILLIN

NOW... Council Accepted



BICILLIN (dibenzylethylenediamine dipenicillin G) is a new penicillin compound. It possesses characteristics which set it apart from older forms of penicillin. Unique is BICILLIN's relative insolubility; its tastelessness; its resistance to gastric degradation; the apparent ease with which patients tolerate it; the stability of its oral forms. BICILLIN indeed opens to view new horizons in antibiotic therapy . . . new applications of penicillin—drug of choice in a wide range of infections.

BICILLIN is available in oral suspension, tablet and injectable forms



Philadelphia 2, Pa.

2 Convenient dosage forms

'Ilotycin,' Crystalline, Tablets
(ERYTHROMYCIN, LILLY)

'Ilotycin Ethyl Carbonate, Crystalline'
(ERYTHROMYCIN ETHYL CARBONATE, LILLY)

Pediatric

- Taste-tested by Lilly's Junior Taste Panel, the flavor of 'Ilotycin Ethyl Carbonate, Crystalline,' Pediatric, is sure to please your young patients—as well as adults who prefer liquid medication.
- Well tolerated—There is minimal gastric irritation, nausea, and diarrhea.
- 'Ilotycin' does not predispose to monilial superinfection or vitamin B complex deficiency.

AVERAGE DOSAGE:
Thirty-pound child, 1 teaspoonful 'Ilotycin Ethyl Carbonate, Crystalline,' Pediatric, every six hours; others, in proportion to weight. Adult, 200 mg. every four to six hours.

SUPPLIED: 'Ilotycin,' Crystalline, in 100-mg. or 200-mg. specially coated tablets. 'Ilotycin Ethyl Carbonate, Crystalline,' Pediatric, is supplied in 60-cc. packages. Each teaspoonful (5 cc.) contains 100 mg. of 'Ilotycin' as the ethyl carbonate.

Lilly

**For successfully
treating the
most common
bacterial infections**



ELI LILLY AND COMPANY • INDIANAPOLIS 6, INDIANA, U.S.A.

ARIZONA MEDICINE

Journal of Arizona Medical Association

VOL. 10, NO. 10



OCTOBER, 1953

Original ARTICLES

INDICATIONS FOR IRRADIATION OF BREAST LESIONS

U. V. Portmann, M.D.
Tucson, Arizona

There are only a few indications for irradiation of benign diseases of the breast. The principle one is for the analgesic effect when there is pain from acute or subacute mastitis, adenosis, or chronic cystitic mastitis.

Tumors of the breast which appear clinically to be benign should be excised because some will prove to be malignant when examined microscopically. Examination should be made immediately, preferably by frozen sections. If cancer is found more radical surgical procedures are indicated at once.

The principle indication for irradiation is when cancer is found. In general about 85 per cent of cases with small movable cancers and no metastases, or stage I, survive five years after operation alone. But 15 per cent die of cancer. This may be because (1) unrecognizable metastases to substernal lymph nodes which may develop especially from inner quadrant tumors, (2) cancer cells may pass through the axillary node group to metastacize in the supraclaviculars or elsewhere, (3) metatases may escape meticulous microscopic examination of axillary nodes, (4) lymph nodes with metastases in them may not have been removed.

Stage II cases have small freely movable tumors with metastases in only a few axillary nodes. The general five year survival rate is reduced to 35 per cent or less by operation alone.

Post-operation roentgen therapy is indicated for stage I and II cases. Local recurrences on the chest wall seldom occur in such cases, but metastases in regional nodes or elsewhere may

develop. Post-operation irradiation should be given to the regional lymph node areas including the substernal groups. It may not be necessary to irradiate the chest wall if there is reason to believe that all of the malignant tissue has been removed. The irradiation may not cure but will increase the five year survival rate in 15 to 20 per cent of cases by destroying some neoplastic cells thereby delaying local growth and remote metastases.

Operations are contra-indicated for stage III and IV cases except occasionally for removal of ulcerations which do not respond to adequate irradiation. Stage IV cases have remote metastases which of course contra-indicates operation. Incidentally all patients to be operated upon for tumors of the breast should have radiographic examinations of the chest, vertebrae and pelvis prior to operation to eliminate the possibility of metastases in these regions.

Irradiation for remote metastases for stage IV cases is purely palliation. Sex hormone therapy may be given concurrently. In older women stilbesterol seems to be best by oral doses of 10 to 30 mg. per day. In younger women testosterone propionate intra-muscularly in doses of 100 mg. three times a week may be given but half this dose often is effective. Also ablation of ovarian functions by roentgen therapy seems to be beneficial in some cases.

The principle indication for irradiation and benefits for cancer of the breast is for stage III cases which are incurable and inoperable. The criteria of incurability should be carefully sought and women with them not subjected to opera-

Presented before the Annual Meeting of the Arizona Medical Association held at Tucson, April, 1953.

tion but treated by irradiation alone. These criteria areas follows:

The Skin

1. Edema (orange or pink skin) of more than slight extent.
2. Ulceration of more than slight extent.
3. Brawny red and inflamed not obviously due to infection.
4. Secondary skin nodules.

The Breast

1. Diffusely edematous.
2. Diffusely infiltrated
3. Multiple secondary tumors.
4. Fixation to chest wall.

Metastases

1. Axillary lymph nodes numerous, extensively involved and fixed.
2. Supraclavicular lymph nodes or edema of arm.
3. Involvement of contralateral breast or lymph nodes.
4. Remote metastases in bones, lungs or other viscera.

Eliminating the presence of remote metastases of Stage IV, the criteria given are those of stage III. An analysis which I made of 136 Stage III cases shows the following results in percentages alive following operation alone and x-ray therapy

alone in yearly periods up to five years.

	No.	1	2	3	4	5
Alive operations only	53	49.5	17.0	6.0	4.0	4.0
Alive x-ray alone	83	83.3	54.1	33.3	20.8	17.0

It will be noted that only half of the patients operated upon survived one year and 4 per cent survived five years. In all probability 4 per cent would have survived as long without treatment. Also the percentage of survivals in each yearly period of patients given roentgen therapy alone is higher than those operated upon and 17 per cent lived five years. This was a worthwhile extension of economic usefulness and life.

CONCLUSIONS

1. Irradiation may relieve pain associated with certain benign disease of the breast.
 2. Small tumors of the breast should be excised, examined microscopically at once and if cancer is found more radical operations are indicated without delay.
 3. Stage I and II cases of cancer of the breast receive postoperation roentgen therapy to delay extension.
 4. The criteria of incurable cancer of the breast have been given.
- Stage III and IV cases should not be operated upon but treated by irradiation alone. Also the use of sex hormones and ablation of ovarian functions may be beneficial.

THE ROLE OF THE NON-SPECIALIST IN THE CARE OF ALLERGIC DISEASE

Samuel M. Feinberg, M.D. *
Chicago, Illinois

Clinical allergy is at the crossroads! Unless we can interest the non-specialist in the care of at least some of the simple allergic problems, the prevention and management of allergic mishaps arising from diagnostic and therapeutic procedures, and cooperation with the allergist in difficult cases, millions of allergic individuals are doomed to unnecessary suffering. The role of the non-specialist in the management of allergic disease should be actually a greater one than that played by the allergist. The purpose of this presentation is to indicate to you the reasons why the non-specialist must deal with allergy, the know-how of importance to him and how this knowledge can be obtained.

*Presented before the annual meeting of the Arizona State Medical Society, at Tucson, April 27, 1953. From the Department of Medicine and the Allergy Clinic, Northwestern University Medical School, Chicago.

WHY THE NON-SPECIALIST MUST DEAL WITH ALLERGY

Prevention and care of man-made allergic manifestations. One of the big problems in medical practice today is the prevention and care of allergic syndromes arising from diagnostic and therapeutic procedures. This phase of medicine was negligible 50 to 75 years ago, when the chief diagnostic instruments were the stethoscope and fever thermometer, and when therapy was limited to blood-letting, calomel and a few other relatively simple drugs. Quinine, aspirin and coal tar derivatives then became part of our armamentarium and thousands of cases of allergy to these drugs ensued. Urticaria is one of the lesser manifestations. Asthma from aspirin and other analgesic drugs is not rare, particu-

larly in those who are subject to chronic asthma. Asthma induced by aspirin is extremely violent and sometimes fatal. In addition to these effects it became evident that some drugs, particularly aminopyrine, were capable of producing hematologic changes, especially granulocytopenia.

Then came the era of arsphenamines, with the concomitant advent of thousands of cases of allergic reactions, many of them serious and some fatal. The use of animal sera for prophylaxis and cure of infections and diseases of intoxication began to demonstrate a train of untoward responses such as serum sickness and the immediate anaphylactic type of reaction, some of which were fatal.

Almost from the very inception of the use of sulfonamides allergic effects to these drugs have been noted. Rashes, "serum-sickness" type of manifestations, occasional anaphylactic type of response, granulocytopenia, aplastic anemia, thrombocytopenia and periarteritis have been among the effects observed.

The newer antibiotics have created a new problem in allergy which must be faced by every practitioner of medicine. Topical applications of penicillin and some other antibiotics to the skin have been largely abandoned because of their great tendency to produce contact sensitization. The delayed type of reaction from penicillin, occurring customarily about ten days after its administration, and consisting mainly of urticaria, angioneurotic edema, arthralgia and fever, still affects 2 to 5 per cent of those receiving the drug. The use of different types of penicillin or the concomitant use of antihistamines has had no appreciable effect on this tendency. More recently it has become apparent that a more dangerous type of penicillin allergy is on the increase. Due to repeated use of the drug, an increasing number of people are acquiring the anaphylactic type of sensitivity. In mild form this may consist of symptoms of urticaria or asthma immediately following administration of the drug. In its more severe form the patient goes into shock and frequently becomes unconscious. In such instances a fatal outcome is not unlikely. There is good reason to believe that the fatalities from such anaphylactic episodes now amount to several score. Such reactions presumably may result from any type of penicillin, and recently it has come to light that the iodide salt of penicillin, penethamate hydri-

dide (neo-penil^R) is particularly likely to produce such effects.

To these man-made allergies one should add those arising from insulin and liver injections, hormones, vitamins, vaccines, contrast media for gallbladder and kidney visualization, and many drugs such as opium derivatives and others. Also to be included are allergic reactions resulting from transfusions. Usually these take the form of urticaria, but other and more serious manifestations may result. They are caused either by the reaction of the donor's allergic blood with a food (or drug?) which the recipient has ingested or by the interreaction of the recipient's allergic serum with an antigen from the donor's blood. Such reactions are, of course, minimized by pooled plasma.

Care of simpler and acute allergic disease. There are about 10 million people with allergy in this country, about half of whom require fairly regular or repeated care and the other half occasional attention. The number of physicians who confine their practice to allergy and who are reasonably well qualified does not exceed one thousand. Ten thousand patients per allergist is an impossible load and it is quite obvious from the standpoint of numbers alone that a large percentage must seek relief from the non-specialist.

It is no more than right that the internist, general practitioner and pediatrician should undertake the care of the simple cases of hay fever, and should be able to manage the patient with the occasional or mild asthma. The rhinologist is expected to prescribe for the patient with an allergic rhinitis or even at times to look for the specific etiologic factor. The acute case of urticaria must, in most instances, be managed by the non-specialist. The successful pediatrician must learn how to handle the problem of infant feeding from the standpoint of the prevention and management of food allergy. Even the surgeon and the obstetrician should recognize that they encounter allergic problems. In addition to the allergy arising from therapeutic and diagnostic procedures, important items include the tendency to postoperative atelectasis in asthmatic patients, the choice of an anesthetic, the effect of pregnancy on the mother's allergy and advice as to the possibilities of inheritance of allergy. It is the family physician on whom the patient must depend when he has an acute allergic illness, whether it be hives or asthma.

Cooperation with the allergist. The allergist requires the cooperation of the non-specialist in the care of many of the allergic patients. Since allergists cannot be planted at distances of several miles apart it is obviously difficult for many patients to come to the specialist for regular and repeated treatments. It is here where the general practitioner, internist, pediatrician or other type of specialist must act as an associate to the allergist. He should carry out the suggestions, advice and give the injections prescribed by the allergist. The non-specialist may have to see that the patient carries out the diets and other suggestions. As a family doctor he is frequently in a position to learn many things about the patient — his habits, emotional factors and the physical environment of his home and occupation — details which may escape the specialist in his isolated consultation room in the heart of a large city. The physician who is closer to the patient may frequently have to prescribe medication indicated for the conditions at the particular moment.

The non-specialist and allergist should regard themselves as true associates in the care of the patient. The allergist should be informed periodically of the progress of the patient, new developments and any other points of interest. He should not be regarded as a glorified laboratory technician who issues a report to the doctor, or even as a casual consultant who after seeing the patient once, has terminated his interest and usefulness. The allergist, on the other hand, must not regard the cooperating physician as a technical assistant or in the capacity of a nurse who is expected only to follow a list of doses and give a few injections to the patient. The allergist must see to it that the cooperating physician is fully informed about the patient's findings, the rationale of his treatment, the composition of the antigens, the goal to be achieved and anything else that the allergist knows about the patient. Above all, each must have confidence in the other. Under these favorable conditions the non-specialist need not fear the loss of practice when he refers a patient to the allergist. Not only can he, in many instances, continue to care for his patient, but even where he cannot or does not wish to do so, the successful outcome increases patient and family goodwill and results in ultimate gain to the referring physician. To the allergist, of course, the conscientious cooperation of the non-specialist is a

matter of absolute necessity. Without him he cannot succeed.

INFORMATION ON ALLERGY OF IMPORTANCE TO NON-SPECIALIST

It is apparent that all of this presupposes that the non-allergist knows something about allergy. Obviously at this point it is not possible to go into detail of what this knowledge should be, but it is appropriate to suggest the main topics of information that should be part of the practitioner's armamentarium.

The causes and manifestations of allergy. The mere recognition of the role of many drugs in the causation of allergic manifestations will do much to minimize the possibility of such occurrences. The knowledge that some gastrointestinal manifestations or that some headaches may be due to food allergy, the realization that pollen and molds cause the majority of seasonal hay fevers, and the willingness to face reality in the asthmatic child and not wait for it to "outgrow" the asthma, are but examples of bits of information which may be ultimately of help to the patient. It's important to know that an allergic cause cannot be found by the honest allergist for every case of asthma or vasomotor rhinitis. It is equally important to be hard-cored about accepting syndromes into the allergic fold just because they are puzzling and do not have an immediate explanation at hand. Chronic fatigue, muscular pains, insomnia, jitteriness, schizophrenia, acne and chronic alcoholism are not allergic manifestations in spite of the claim of some extremists to the contrary. Neither has it been proven that refined food products, such as traces of starch, purified dextrose or cane sugar carry the allergenicity of their food of origin.

Principles of diagnosis. Much can be learned about the allergic patient from history-taking. The history in some instances is far more important than any other diagnostic procedure. The closer the physician is to the patient the more informative the history may be. This is then to be regarded as a major instrument of the non-specialist. The history may suggest the presence of certain complications or it may give a clue to the diagnosis of the specific cause. And frequently it may warn the physician what medication not to give.

The elements of skin testing should be known whether the physician does the tests himself or depends on a report from someone else. For atopic disease (asthma, hay fever, eczema, etc.)

the appropriate skin test is either the scratch or intradermal procedure. The scratch test is safer, more often clinically significant and less painful. The intradermal test is more sensitive but at the same time lends to a greater incidence of non-significant reactions, it is more painful and may be hazardous. Many allergists use the intradermal to supplement the scratch method. For the non-specialist the scratch technique is preferable. A positive test consists of an urticarial wheal, with erythema and itching occurring within 5 to 20 minutes after the application. The patch test is employed in contact dermatitis. It is read in 24 to 48 hours. A positive reaction consists of itching, erythema and possibly vesiculation.

The interpretation of the skin test is a major concern of the non-specialist. To assume that a positive skin test solves the cause of the patient's complaint is to open the door to endless error and much injustice to the patient. A positive skin test may also mean that the antigen is related to the one which causes the allergy, that it's the cause of an allergy of which the patient does not complain or that the clinical manifestation may not show itself months or years later. One must be aware that certain substances, such as histamine, morphine, codeine and acetylcholine, give a positive whealing reaction in everyone. Many solutions or preparations used for testing are irritating and the resultant responses are false and misleading. Dermographism is a frequent source of false reactions.

Conversely, a negative test must not be taken at its face value. In some allergic conditions such as urticaria and headaches, the tests are usually negative even when the causative allergic factor is known. In "serum sickness" type of reactions, consisting of urticaria, arthralgia and fever, occurring several days after the administration of serum, penicillin and some other therapeutic substances, the skin test is almost always negative. Such manifestations cannot be forecast by a skin test prior to the administration of the drug. On the other hand, the anaphylactic or immediate type of reaction to penicillin or serum is usually associated with a positive whealing skin test. In view of the growing hazards from penicillin, it is advisable that the administration of this drug in those whose have had it previously be preceded by a scratch test. If the latter is negative an intradermal test (10

to 100 units to 1 cc) may be used. If the test is positive the use of penicillin is hazardous.

In spite of their capacity to produce allergy most drugs do not give positive skin tests. This is particularly true of aspirin, aminopyrine, phenacetin and related drugs. Aspirin may produce urticaria or such violent asthma that at times a single tablet has resulted fatally. The informed physician will not be fooled by the misinformed laboratory which issues a report that the patient has a negative skin test to aspirin and is, therefore, not allergic to that drug.

Principles of treatment. Since the non-specialist must administer most of the palliative treatment for allergy he must be cognizant of its major principles. Adrenalin is the best remedy for an allergic emergency, such as an anaphylactic reaction or acute asthma. But it is of little use in serum sickness, where antihistamines are preferable. The antihistamines have limited value, being useful particularly in the itching dermatoses and in allergic rhinitis; their effect in asthma is negligible. The iodides are still an excellent remedy in persistent asthma. Aerosols of epinephrine or isopropyl epinephrine (isuprel or norisodrine) are useful adjuncts in the treatment of asthma. ACTH and cortisone are valuable remedies in some cases of acute status asthmaticus and for the maintenance therapy of occasional persistent allergies, such as some cases of intractable asthma. They certainly should not be used as a first or routine measure, as a substitute for diagnosis or as a substitute for other more simple and less hazardous symptomatic remedies.

There is nothing more dramatic and more complete in the therapy of chronic disease than the relief of a chronic allergy by the removal of the specific cause. This should be the objective in the treatment of allergy whenever possible. To this end the non-allergist must learn a few elementary principles. The basic procedures in formulating a diet, the procedures for avoidance of house dust and the substitution of safe cosmetics are among such steps.

It is in the realm of desensitization therapy that principles of good practice are frequently abandoned. Desensitization should only be given for inhalant or contactant allergy. Inhalants should be used only if the cause is proven and avoidable. Pollen from plants which are insect

pollinated, dander of a dog which can be more easily removed from the home, or pollen from timothy which is hundreds of miles away do not call for desensitization therapy. For the most part, pollen, molds and house dust are the chief antigens used in desensitization, while occasionally the use of special, occupational or other unavoidable inhalants may be indicated. There must be a rational causal relationship before such therapy is employed. The principle of the use of small, harmless doses, of gradually increasing amounts should be followed. Conservatism should be the rule and reactions should be avoided. Desensitization therapy requires care and certain precautions but mastery of its use can be accomplished by the discerning physician.

Most physicians have had little if any opportunity to learn the principles of allergy when they were medical students. It is true that the curricula of most medical schools are giving increasing emphasis to allergy. Nevertheless, the practicing physician must discipline himself to acquire knowledge and experience from sources other than his undergraduate training. Short postgraduate courses are frequently available

and helpful, providing that the physician does not delude himself into thinking that a course of a few days makes him a specialist. Allergy programs are frequently offered, such as by symposia at the state or county meetings or at the Allergy Session of the A.M.A. Regional and national allergy societies also offer programs from which the non-specialist can obtain considerable benefit. And last, but not least, there remains the valuable training derived from careful observation of patients, discriminative reading and study and the free exchange of ideas in the doctors' lounge in the hospital.

SUMMARY

The care of allergic patients requires both the allergist and non-allergist. The allergist is needed for the diagnosis and therapeutic guidance of the difficult cases. The non-allergist must handle the numerous simple cases of allergy and must learn how to prevent and manage the allergy arising from drugs. To the allergist and to the allergic patient the non-specialist is indispensable in the cooperation required to carry out management outlined by the specialist. With reasonable effort it is possible for the non-specialist to equip himself for this task and duty.

WHY REPORT COMMUNICABLE DISEASES?

Clarence G. Salsbury, M.D.

James B. Swayne, M.A.

Phoenix, Arizona

50 Years of Reporting

1953 marks the 50th Anniversary of compulsory reporting of contagious diseases in Arizona. The legislative act of 1903 creating the Territorial Board of Health established the legal framework for reporting. Since this original legislative enactment is still a part of State law, it is of more than historic interest:

68-303. *Report of Contagious Disease*—If it shall come to the knowledge of any person that a contagious, epidemic, or infectious disease exists he shall immediately report in writing to the Board of Health having jurisdiction the name and place of residence, if known, of every person affected with such disease and if he is the attending physician he shall report not less than twice in each week the condition of the person afflicted and the state of such disease. (Laws 1903, Ch. 65).

Federal legislation on collection of morbidity data preceded the Arizona enactment by about

25 years. In 1878, Congress authorized the U. S. Public Health Service to collect data for use in connection with quarantine measures against such pestilential diseases as cholera, smallpox, plague and yellow fever. (1) In 1893, another law was enacted authorizing the U. S. Public Health Service to collect weekly data from states and municipalities. In 1902, the Surgeon General was authorized to provide forms for the collection, compilation and publication of communicable disease data. Throughout much of its existence in Arizona, the program has been a cooperative federal-state-local venture. It has been modified somewhat from time to time to reflect changing conditions, but essentially has always involved the notification of health authorities of the names and addresses of persons with communicable disease.

Recent Developments Nationally

The latest major change nationally was made in 1951 when a committee of epidemiologists and

statisticians met at the Communicable Disease Center of the Public Health Service and viewed the list of reportable diseases to determine those which remain of national interest and to consider methods of improving the mechanisms for obtaining data. Following this meeting a revised national list of diseases was adopted and a procedure set up for publication weekly of brief narrative reports on epidemics and occurrences of rare disease — these reports to be channelled from local health authorities and epidemiologists through state health officers to the Public Health Service. This latter step was taken to supplement the existing system of weekly reports. One of the factors behind adoption of this mechanism for exchange of information was the possibility of biological warfare with the resultant need for a fast system of epidemic intelligence. By rapid amassing of information from all over the country, possibly significant events can be quickly screened and early control measures taken when indicated.

Recent Developments in Arizona

A radical change in the system of collecting communicable disease data was made in Arizona in January 1953. The former system was based on weekly numerical postcard reports from physicians supplemented by individual case reports on a fairly long list of diseases including tuberculosis, syphilis, gonorrhea, poliomyelitis, brucellosis, typhoid, etc. Several different types of case report forms were used — the type used depending on the disease.

Weekly postcard reports on the number of cases were eliminated under the new system. A single case card was adopted which can be used for any reportable disease. A brief description of the reporting system together with a cut of the report card and envelope is presented in the accompanying exhibit, "How to Report Communicable Diseases."

Value of Reports

Discussions of the value of any statistical reporting program are apt to fall into generalities which neither inform nor motivate desirable action. It is hard to avoid generalization and still retain brevity in a discussion of the value of communicable disease reports. A truly adequate statement must consider separately the special characteristics of each disease — the nature of available prevention and control measures, how the program actually operates, and the status of scientific research and possible need

for information on incidence to help in developing new preventive measures.

Such a statement on value or use of communicable disease reports must also consider the needs of a wide variety of users. A list of recent special requests for morbidity data received by the State Health Department includes such voluntary health agencies as the National Foundation for Infantile Paralysis, the National Tuberculosis Association, various government agencies, newspapers, magazine feature writers, commercial pharmaceutical firms, research organizations, and schools of public health. The data receives an even wider distribution by the U. S. Public Health Service which regularly compiles and publishes data received from each state and territory.

The uses made of communicable disease data are constantly changing and it is not far-fetched to believe that the next 12 months may see some new use arise which cannot be anticipated today. The evolution and mutations which have occurred will be evident from the discussion which follows.

Disease Prevention and Control

The original purpose of communicable disease reporting was to enable officials to enforce quarantine measures. Today more emphasis is placed on preventive measures and less dependence on quarantine. This shift in emphasis, however, has not changed the need for case reporting.

There is a group of virulent diseases on which reports are required such as smallpox, diphtheria, rabies, malaria, and typhoid fever which have been virtually eliminated. The conquest of these important contagious and infectious diseases is one of the major triumphs of science, medicine, and public health. The great gains made in the control of these diseases, however, cannot be retained without continuance of control measures. Knowledge of even a single occurrence of any of these diseases is important. The report rings the alarm which indicates something has gone wrong with control measures; it makes it possible to bring community forces into action so that defects in the control program can be eliminated and further spread of the disease prevented.

There still remain a number of contagious and infectious diseases for which control measures are available but which continue to be quite common. Control programs such as those for

tuberculosis and venereal disease cannot operate effectively without case reports. By way of illustration, contacts cannot be followed up and brought to treatment unless physicians report the cases which they diagnose. Such follow-up, it should be stressed, is made only with the express permission of the physician.

Gamma Globulin Allocation

A new use of communicable disease incidence data which could not have been anticipated even a year ago is for allocating the limited supplies of gamma globulin. With the recent discovery that gamma globulin appears to afford a measure of temporary protection against polio by reducing the severity of paralysis, the potential demands on the limited supplies available require some fair system of allocation which will assure that available supplies are utilized where the need is greatest. Accordingly, each state will receive an allocation for polio based on the median number of cases reported during the period 1947-1951 inclusive. Emergency allocations for polio will then be made where severe outbreaks occur — the evidence of severity being officially reported cases. Parenthetically, it should be said that during the last polio season (1952) the bulk of reports were received from four to eight weeks after onset, and then in many instances only after follow-up of leads received from sources other than the attending physician.

To improve allocation of gamma globulin, the U. S. Public Health Service has requested that reports on polio cases be provisionally classified as paralytic or non-paralytic. This classification is necessitated in part by the high degree of uncertainty in the diagnosis of non-paralytic cases and the need to have knowledge of severe outbreaks so that officials responsible for allocation of supplies can more accurately determine the areas where need for additional supplies is greatest. The State Health Department joins the Public Health Service in requesting that physicians tentatively classify polio cases as paralytic or non-paralytic when submitting the original report on a case.

The danger of shortage of gamma globulin demands extreme care in utilizing available supplies for measles and infectious hepatitis. Accordingly, the State Health Department has adopted the policy that supplies will be allocated only where reported data show that need exists. This policy is influenced somewhat by the knowledge that if a really acute shortage

occurs requests by the state for supplementary supplies for use with measles and infectious hepatitis will probably be met only if factual data are available to support the request.

Limitations of Mortality Data

In the past, considerable reliance could be placed on mortality data from death certificates to provide an index of the incidence or prevalence of certain communicable diseases. However, with the rapid forward strides of science and medical practice, mortality from most of the communicable diseases has been or is rapidly being virtually eliminated. For some diseases, however, the decline in mortality has not been accompanied by a proportional decline in incidence. This is true of tuberculosis where conclusions as to progress made in control may be quite false if based on mortality data alone. For this reason, it is more important than ever to improve tuberculosis reporting so that it will be a sensitive indicator of problems and progress toward the goal of eliminating the disease as a public health problem.

Inadequacy of Numerical Reports

For many years prior to 1953, Arizona collected communicable disease data through a system which made use of weekly numerical postcard reports from physicians, supplemented by individual case reports for a number of diseases. The elimination of the weekly postcard reports in January 1953 can in no sense be considered an Arizona innovation. A national study in 1950 of morbidity reporting procedures used by the various states showed only three were using weekly reports from physicians, giving totals by disease. The remaining states all depended solely on individual case reports which gave as a minimum the name and address of the patient. (2) As far back as 1936, Dr. Carl Buck, then Field Director of the American Public Health Association made the following comment in a survey of Public Health in Arizona: (3)

"The present system of reporting on a card which merely indicates the number of cases of certain diseases and gives nothing concerning their specific location is almost useless and should be immediately abandoned. Cards should be supplied to physicians for reporting communicable diseases and such card should provide for all necessary pertinent data concerning that case, particularly name, age and specific address."

It will have been noted that the original legislation which required reports of contagious dis-

eases required an individual case report giving the "name and place of residence, if known, of every person affected with such disease". Our research does not disclose when or why a shift to numerical totals was made nor do we know why Dr. Buck's recommendation for elimination of weekly postcard reports was not carried out. We believe that the principal factor may have been a reliance on the weekly postcard to obtain clues for follow-up to secure individual case reports.

The weekly numerical postcard report was simple to complete, but experience showed that it invited estimates and errors. Some physicians based their weekly reports on memory rather than referring to office records before preparing their reports. Many saved the reports and sent several weeks' reports in at one time. Follow-up on several "tularemia" outbreaks showed that the reporting physician had intended to report "syphilis" — the preceding item on the alphabetical list of diseases. In some instances the same tuberculosis cases were reported week after week. This was probably due to failure of an office assistant to understand that only new cases should be reported. These types of errors were discovered because they involved diseases for which case reports were requested. The amount of error in tabulation of diseases for which case reports were not requested can only be guessed at.

Case Reports Essential

Case reports giving name, address, and certain basic facts such as age, sex, race, and date of onset are basic to good morbidity reporting. Accuracy is improved by obtaining names, since duplicates and previously reported cases can be eliminated. Efforts to improve reporting can be extended because health departments can check on reporting of cases which came to their knowledge from sources other than physicians.

Addresses are essential for statistical allocation of cases to specific counties, cities or smaller areas within cities. The seasonal distribution of a disease is only available when date of onset is reported. Data on geographical distribution correlated with date of onset is basic to many epidemiologic studies. The age, sex, and race distribution is often fundamental for research.

Apart from statistical considerations, names and addresses must be reported for any disease where quarantine or follow-up of suspects is a part of the control program or if the physician wishes to refer cases for public health nursing

services. The case report is also essential for epidemiological follow-up on cases to determine factors responsible for origin and spread of an outbreak.

The foregoing is by no means a complete resume of the interests of the users of communicable disease data, but it suggests the reasons which impelled inclusion of the minimum core of basic information on the case card. Although the demands for other types of information are great, a real effort was made to include a minimum number of items because of the demands on the physician's time and the desire not to make the system unworkable because of the number of items required.

The shift from weekly postcard reports to complete reliance on case reports required two basic essentials. First, a simple case card which could be used for any disease. Exploration of the experience of other states showed that California had recently adopted a single case card. Using the California card as a basic pattern, a few modifications were made to adapt the card to Arizona's needs. The second essential was a reduction in the list of diseases to the very minimum consistent with a good health program. Following this review, it was decided not to require case reports on influenza, chickenpox, german measles, impetigo, mumps, and ringworm.

Influenza is a disease of considerable public health significance because of the disability and economic loss it causes. It was not included on the list because experience has shown that reporting is notoriously inadequate due to the lack of use of specific diagnostic procedures, similarity in symptoms to other common diseases, and the vast number of cases which occur in epidemic periods. New techniques are being developed on an international and national level in an attempt to obtain the data needed for research and control.⁽⁴⁾ Although case reports are not required for influenza or the various childhood diseases listed in the preceding paragraph, physicians are requested to report unusually high incidence.

Criticism has been received because measles was included on the list of diseases. We have already seen the value of measles information in the allocation of gamma globulin, and have noted the difficulties encountered with numeric reports where names and addresses are not given. The authors recognize, however, that the high incidence of measles during certain periods

creates a real problem. This matter of securing data on incidence of measles will be reviewed again and an effort made to arrive at a satisfactory solution. Comments and suggestions from all interested persons on this matter or on any aspect of the reporting program will be appreciated.

Reporting Lacks

The compiled data on reportable disease suffer from all or at least nearly all the disabilities that a statistical series can have and still survive. These disabilities include gross failures to report, bias due to difficulties of diagnosis, and human carelessness. All these factors are inter-related, but perhaps the greatest single fault is incompleteness. Obviously cases which do not seek medical treatment cannot be reported, but there are tremendous gaps even for those cases which do seek treatment. When the weekly postcard system was used there were relatively few weeks when even 50 percent of the physicians submitted the postcard report. For tuberculosis, where there is a fairly extensive case finding program, almost two out of three deaths which occurred in Arizona in 1952 had never been reported as a case.

Resistance to communicable disease reporting, apart from the question of case reports on measles, seems largely to center on two objections. First is the desire to protect the confidential nature of the doctor-patient relationship. The most pertinent comment here is that the local and state health departments as custodians of the records take all reasonable means to protect the privacy of any individual whether he be the patient of a private physician or of a public clinic. Since reporting is a legal requirement which recognizes the primacy of the public interest, there seems little moral or ethical grounds for objection as long as the records are not abused. The second principal objection to reporting communicable diseases is the amount of work required of physicians who are greatly overburdened. Most physicians have office assistants to relieve them of routine work, and preparation of the simple case report card is routine. The physician's signature is not required on the report card although name of the physician is requested. The physician reader may forgive the sly suggestion that case reporting would be improved if preparation of cards were turned over to office assistants 100 percent.

With all the shortcomings of the case report-

HOW TO REPORT COMMUNICABLE DISEASES

Each physician has a booklet containing 25 Confidential Morbidity Report cards. The list of reportable diseases is on the inside back cover.

COMPLETE THIS CARD THE DAY YOU DIAGNOSE THE CASE. (YOUR OFFICE ASSISTANT MAY DO IT FOR YOU.)

Report diseases listed on inside back cover.

MAIL THE REPORT IMMEDIATELY IN ONE OF THE PRE-ADDRESSED POSTAGE-FREE ENVELOPES.

WHEN YOU NEED MORE CARDS OR ENVELOPES FILL IN THE SPACE ON THE BACK OF THE CARD.

For information write to:
Division of Vital Statistics
Arizona State Department of Health
Phoenix, Arizona

ing program it is fair to ask, "Why waste time and money compiling figures which are incomplete and inaccurate?" One answer is that the reported statistics provide basic clues for estimating trends in prevalence and the size of a disease problem. If the user is aware of the limitations of the data and makes allowance for the various factors which affect their interpretation, he may derive useful information which is available from no other source. Without case reporting, other more expensive and less satisfactory means would be required for case finding and for developing the basic data needed for planning and research. The greater the completeness of reporting the less waste there is in operation of public health programs since less effort must be directed toward case finding efforts. The physician who regularly and promptly submits reports not only has the satisfaction of observing the law, but makes an important contribution to public health.

1. National Morbidity Reporting Manual of Procedures; National Office of Vital Statistics, Public Health Service, Washington, D. C.
2. Morbidity Reporting Procedures in State, Territorial and Independent City Health Departments; National Office of Vital Statistics, Public Health Service, Washington, D. C. (January 1951).
3. A Survey of Public Health in Arizona; Carl E. Buck, Dr. P. H., 1936.
4. Influenza Study Program in the United States; Public Health Reports, Vol. 67, No. 12.

PHOENIX *Clinical* CLUB

MASSACHUSETTS GENERAL HOSPITAL CASE RECORD NO. 24

The Case History in this discussion is selected from the Case Records of the Massachusetts General Hospital, and reprinted from the New England Journal of Medicine. The discussant under Differential Diagnosis is a member of the staff of the Massachusetts General Hospital. The other discussants are members of the Phoenix Clinical Club.

A six-and-one-half-year-old boy was admitted to the hospital because of rapid weight gain, growth of pubic hair and enlargement of the penis.

On routine examination nine months before admission, he had been found in good health, with normal development, weighing 44 pounds. The blood pressure was 100 systolic, 60 diastolic. The genitalia were noted as normal. Approximately six months before admission, however, he began to gain weight rapidly and to develop pubic hair and penile enlargement. He was otherwise completely asymptomatic and had been very energetic and active, with good muscular strength. He had a good appetite without special preferences, although he ate a great deal of meat and potatoes. There was no special craving for salt or sugar. There were no voice changes, development of axillary hair, acne or other skin changes.

The past history revealed that he was born in breech presentation at term and weighed 6 pounds, 10 ounces. Weight gain and development were normal.

A review of the systems was negative except for an episode of "abscessed ears" and tonsillectomy eighteen months previously for frequent upper respiratory infections. He rarely wet the bed, and there were no other genitourinary symptoms.

Physical examination showed a well developed boy. The weight was 65½ pounds, and the height 46 inches. He seemed very plump, with prominent abdomen and hips. The neck and cheeks were fat. Examination of the head, neck and chest was negative. The heart was slightly enlarged to the left. The rate was normal, and there was a Grade II pulmonic systolic murmur. An electrocardiogram gave a normal tracing. The abdomen was plump and full, but no masses

were palpable. There was definite, kinky pubic hair. The penis measured 7.0 cm. in length and 6.3 cm. in circumference. The testes were small and measured 1.9 cm. in cross section. The skin was clear, and there was no acne or unusual pigmentation. There was no axillary hair. Neurologic examination was negative.

The temperature was 99.2 F. the pulse 100, and the respirations 22. The blood pressure was 132 systolic, 96 diastolic.

The urine had a specific gravity of 1.022. The tests for sugar and bile were negative. There were rare red cells and white cells in the sediment. The blood hemoglobin was 14 gm., and the white-cell count 7700, with 78 per cent neutrophils, 14 per cent lymphocytes and 8 per cent monocytes. The eosinophil count was 112 cells per cubic millimeter. The serum chloride was 106 milliequiv., the sodium 140.4 milliequiv. per liter. The nonprotein nitrogen was 17 mg., and the fasting blood sugar 86 mg. per 100 cc. (the blood sugar of a specimen taken two hours after breakfast was 97 mg). A sample of sweat contained more potassium and less chloride per liter than normal. The urinary 11-17 oxycorticosteroid was 0.2 mg. per 24 hours. The 17-ketosteroids were 6.6 mg. per 24 hours.

X-ray films of the right hand and wrist indicated a bone age of seven years. Films of the skull failed to show any significant changes; the bones were of normal texture and density. The pituitary fossa was intact. Chest films showed clear lung fields. The heart was at the upper limit of normal size. The aorta was prominent. Anterior and posterior views of the abdomen showed normal appearance of the left kidney, spleen and liver. The right kidney was obscured by intestinal contents. There was no evidence of abdominal tumor.

On the fourth hospital day an operation was performed.

DR. LOUIS B. BALDWIN

This six year old boy had precocious sexual development over a period of six months with gain in weight, growth of pubic hair and enlargement of the penis without proportional enlargement of the testicles. There were none of the other evidences of puberty, such as, change in voice, the development of axillary hair and acne.

It is further to be noted that he appeared perfectly well but that the heart was slightly enlarged and that he had an elevated blood pressure for his age.

The laboratory studies were all normal with the exceptions of a possibly slightly lowered blood potassium and the statement that, "a sample of sweat contained more potassium and less chloride per liter than normal."

This brings us to a consideration of the various causes of precocious sexual development in the male.

Pineal tumors may produce precocious sexual development. As a matter of fact precocity is present only exceptionally in tumors of the pineal, and it is now felt that pubertas praecox in pineal tumors is due to pressure exerted by these neoplasms on the contiguous hypothalamic centers and not to a hormone elaborated by the pineal body. This condition seems untenable in our case because of the absence of symptoms of increased intracranial pressure with headache, vomiting, and failing of vision which are almost invariably the first symptoms noted. On the other hand the most important character distinguishing these cases from suprarenal cortex or anterior lobe pituitary disorders is the absence of hypertrichosis. This circumstance isolates this specific hypergenitalism from those related to other ductless glandular dyscrasias. It is interesting to note that Weinberger and Grant collected 17 cases of precocious puberty of whom 3 were females. In all these 17 instances there was disease of the hypothalamus with no evidence of pineal, ovarian, or adrenal pathology. Mid brain tumors were the most common findings associated with this clinical syndrome, although in isolated instances diffuse mid brain inflammatory lesions, such as, encephalitis and non specific inflammations of the brain as well as degenerative encephalopathies were the only abnormalities noted. In these hypothalamic cases there is somatic growth, enlargement of the genitalia with active libido and spermatogenesis.

Although no pituitary tumor has been known to produce precocious puberty or true virilization as its sole manifestation relatively minor virilization symptoms have been observed in association with acromegaly. In pituitary basophilism there may also be precocious puberty but usually associated with the metabolic disturbances characteristic of Cushing's syndrome. It is likely that the virilizing manifestations both in

acromegaly and pituitary basophilism are mediated through the adrenal cortex. Experimentally it is entirely clear that the pituitary influences the size of the adrenal cortex considerably. Smith demonstrated that experimental hypophysectomy caused atrophy of the adrenal cortex.

The normal films of the skull with the pituitary fossa intact does not rule out a basophilic adenoma as these are small tumors, but the absence of symptoms pointing to either acromegaly or pituitary basophilism rule out disease of the pituitary gland to account for the precocious puberty in this case, as does the normal blood eosinophile count.

Malignant tumors of the thymus have rarely been associated with precocious puberty. Three such cases were originally reported by Layton, Turnbull, and Bratton and in each instance marked adrenal cortical hyperplasia was found, in addition to a carcinoma of the thymus. The syndrome produced in these patient is indistinguishable from true Cushing's syndrome.

Neoplasms of the interstitial cells of the testis, the so called Leydig's cell tumors are very rare. Only 40 cases have been reported in the literature by the end of 1949. Of these 11 have been in children. Nation, Edmondson, and Hammack found that 21 collected cases were benign and 5 malignant. Malignant tumors have only been observed in adults. They are characteristically small and in several instances they escaped recognition by the patient. In some cases even the physician could not be certain of the existence of a tumor in the testicle. In this event reliance must be placed on the consistency of the testis as compared to that of the other one. It is usually encapsulated.

Clinical manifestations of an endocrine type were present in one third of the reported cases of Leydig's cell tumor. The majority of these were boys with precocious sexual and somatic development. It is important to emphasize that this is a pseudosexual precocity. Since spermatogenesis is absent. The endocrine effects of Leydig's cell tumors are best studied in the afflicted child. Eleven cases in children have been reported in the literature, and it is important to realize that all — presented definite evidence of premature androgenic stimulation, characterized by precocious appearance of puberty. In some, precocious somatic development was also present. The tumor was benign in every case and was accompanied by gynecomas-

tia in one instances. In all cases developmental changes were first noted about the fifth year of life. After orchidectomy regression of sexual precocity occurred completely in two, partially in three and not at all in the remaining cases. Reports of hormone essays in patients with Leydig's cell tumors were very meagre. Urinary gonadotrophin and estrogen have been quantitated in only one case by Masson and regarded as slightly elevated. The urinary 17-ketosteroids were greatly increased in this same case.

Adrenal cortical hyperfunction in the prepubertal period is more common in females than in males. During this period the underlying lesion is usually a tumor, and most frequently a malignant one. The metabolic abnormalities of Cushing's syndrome are often seen in association with adrenal cortical hyperfunction in the prepubertal group and generally are associated with mild virilizing manifestations. The question always arises if we are dealing with a tumor of the adrenal cortex, or one of the other conditions already discussed. In the female arrhenoblastomas and other tumors of the ovary have to be considered as well as such rare condition as hyperostosis frontalis interna or polyostotic fibrous dysplasia (Albright's Syndrome).

Constitutional precocious puberty is far more common in females than males. The importance of this condition lies in its differentiation from the pathologic causes of true or pseudosexual precocity. There are no statistics as to the actual incidence of this physiologic syndrome, but it undoubtedly is one of the most common causes of isosexual precocity. In the male there is enlargement of the genitalia and testes, spermatogenesis as well as skeletal precocity.

In the attempt to make an accurate diagnosis we wonder whether we have an instance of over activity of the adrenal cortex due either to simple hyperplasia or to a malignant tumor. In our case the important laboratory tests were made and the only possible abnormalities were a slightly low blood potassium and an elevation of the potassium in the sweat with a lowered sweat chloride. These findings in addition to the body contour, which showed moderate obesity with prominent abdomen and hips, and the slightly elevated blood pressure might indicate a questionable Cushing's syndrome such as occurs in adrenal cortical hyperfunction. But there was no palpable abdominal mass. For the study to be complete there should have been pyelo-

graphic studies and possibly perirenal insufflation for attempted visualization of an adrenal tumor.

It would seem logical to assume that an exploratory operation was performed to find a possible adrenal tumor. In hyperfunction of the adrenals in view of the relationship of the adrenal cortex to electrolyte metabolism, notably that of sodium, chloride and potassium one might anticipate that some alterations in the values of these ions might be encountered in patients with cortical hyperfunction. But Willson, Power, and Kepler, in a series of more than 30 cases of Cushing's syndrome 13 of which were due to adrenal cortical tumors, found only three cases in which there were marked changes in the electrolyte pattern of the blood. In regard to the steroids, it is well established that normal men and women excrete both androgenic and estrogenic compounds in the urine. These compounds, however, have their origin not only in the gonads but also in the adrenals. To date, 28 steroids have been isolated from the adrenal. The normal average daily urinary excretion of androgen in the male adult as 17-Ketosteroids is 6.3 to 6.8 mgm. and in the female 4.2 to 5.6 mgm.

The normal values for urinary oxycorticosteroids is from 0.12 to 0.34 mgm. in 24 hours. The values for urinary excretion of 17-Ketosteroids in the various diseases producing sexual precocity in the male are as follows:

Hypothalamus (Pineal) Normal to +
Testes:

- a. Adrenal rests Normal to +
- b. Leydig's cell Normal to + + + +

Adrenal Cortex:

- a. Hyperplasia + to + + + +
- b. Tumor + to + + + +

From this brief review of the possibilities I feel that we must exclude a Leydig cell tumor because of the size of the testes, pituitary basophilism from the X-ray and laboratory studies, involvement of the hypothalamus because of the negative findings of brain pathology. This leaves adrenal cortical tumor or hyperplasia of the adrenal cortex or constitutional precocity which is rare in the male.

Because of the very slightly elevated Ketosteroids for children one hesitates to make a diagnosis of adrenal cortical tumor. But nevertheless this is my diagnosis:

DIAGNOSIS:

Tumor of suprarenal (probably malignant).

Very poor other possibilities:

Hyperplasia of adrenal cortex.

Leydig's cell tumor.

DR. A. J. FILLMORE

This youngster was apparently well up to about six years of age, at which time it was noted that he began to gain weight rapidly; he developed pubic hair and unusual growth of the penis, all other developmental characteristics and manifestations were normal. Such rapid changes as this boy demonstrated must have been alarming, even to a very proud father.

There are a few other physical findings suggestive of an endocrine disturbance: A blood pressure of 132/96 is high for so young a child. And he had begun to show unusual prominence of the abdomen and the hips. His testes were reported as small.

Apparently there is an over production of the masculinizing hormones and the more common causes for an over production of these are primary hyperplasia of the adrenal cortex or an adrenal cortical tumor.

These disturbances are not common, but are much more common in the female than in the male. It is likely the famous case of Christine Jorgensen, recently reported in the press was such a case.

The principal metabolic actions of hormones, or steroids of the adrenal cortex have been defined as: (1) For the maintenance of electrolyte balance. (2) Regulation of carbohydrate metabolism and (3) Promotion of growth and the development of secondary sex characteristics.

The electrolyte regulating steroid, 11 desoxycorticosterone which is similar to, if not identical with, the synthetic steroid, desoxycorticosterone—causes retention of sodium; chloride; and water, and increases the urinary excretion of potassium and phosphorus.

There are two distinct syndromes attributed to hyperadreno corticisin whether from hyperplasia or tumor. 1st: the adrenogenital syndrome and Cushing's syndrome. It is believed that two different hormones or groups of hormones are responsible for the two syndromes. In order to keep the discussion clear the term

Cushing's syndrome refers to changes caused by the adrenal cortical hyperplasia and tumors. While the term *Cushing's Disease* refers to those cases in which basophilic adenomas of the pituitary are present and seem to be responsible for the hormonal changes. When these are absent and the clinical features of the disease are present, the term *Cushing's Syndromes* is applied.

The regulation of carbohydrate metabolism was not disturbed in this youngster as is shown by the fasting blood sugar of 86 mg.% and a non-fasting specimen of 97 mg.%. The process of formation of carbohydrates from the Amino-Acids, glyconeogenesis is governed by the carbohydrate-regulating steroids, of which cortisone, or Compound E is the most widely known.

The most striking changes noted in this young patient are those having to do with the steroids which promotes growth and the development of secondary sex characteristics. And in most cases in males with over activity of the adrenal cortex the primary disturbance consists in excessive secretion of an androgenic hormone. This syndrome of overproduction of the masculinizing steroid varies with the sex and age of the child. The early signs are: enlargement of the penis; scrotum and prostate; the appearance of pubic hair and sometimes facial hair; development of acne and a deep voice. Also a definite increase in height and weight. Our patient certainly had a remarkable growth in his genitals during the interval of nine months, at the beginning of which a routine examination was stated to have revealed no abnormal development. And he had a 20 pound gain in weight during the same period.

It is stated this abnormal growth is not a true sexual precocity because spermatogenesis does not take place. In precocity of hypothalamic origin both penis and testes are enlarged. In sexual precocity of constitutional hypothalamic origin, the urinary 17-Ketosteroids rise only to adolescent or adult levels while in the patient with the adrenogenital syndrome they may be far in excess for an adult. In one case of a negro girl with a tumor of the adrenal cortex she excreted 200 mg. per 24 hour before the tumor was removed. After removal the steroid urinary excretion dropped to 2 mg. per 24 hour. Our patient is reported to have excreted 6.6 mg. in 24 hour, which is 3 to 4 times higher than normal for the average of his age. The increase

in excretion of 17 Ketosteroids is the most important laboratory feature of this syndrome and in male infants and young children it is necessary for a diagnosis of an adrenogenital syndrome.

One of the most important functions of the adrenocortical secretions is the control of the electrolytes. It is generally agreed the major action of the electrolyte controlling hormone is on the kidney, decreasing the excretion of sodium and increasing that of potassium.

There was no apparent disturbance in the electrolyte equilibrium in our patient as it is reported he had a serum chloride of 103 Milliequiv. per liter; and sodium 140 Milliequiv. per liter both of which are within normal limits. The serum Carbon dioxide was not abnormal.

X-ray studies of bones of hand and wrist, and of the skull gave no evidence of disturbance of growth or bony metabolism.

Also x-ray studies of the chest and of the abdomen were not revealing.

It is my opinion that this youngster whose illness was confined to a nine month interval during which time he demonstrated abnormal growth in body height and weight and also abnormal genital development and growth of pubic hair, along with a definite elevation in 17-Ketosteroids excretion, he had an acquired, or post natal Adrenogenital Syndrome. Probably due to a malignant *adrenocortical tumor*. Had the disturbance been due to adreno-cortical hyperplasia the abnormal growth and development of the genitals would have been earlier manifested.

DIFFERENTIAL DIAGNOSIS

Dr. Fuller Albright: This seems like a straightforward case, but one never knows in these exercises.

We know that this patient was producing too much androgen. The rapid somatic growth and the early development of the genitalia are evidence of this. Besides, the 17-ketosteroid excretion was 6.6 mg. per 24 hours, which is very high for a child of this age (normal value, 0 mg.). There are two tissues in the body that produce androgen: the reticular cells of the adrenal cortex and the Leydig cells of the testes. We therefore expect to find some disorder in one of these two tissues.

I will first discuss the reticular cells of the adrenal cortex. The first possibility I would consider is a tumor, benign or malignant, of the adrenal cortex. Everything in this case is com-

patible with such a diagnosis. The rather short duration and rapid progress of the symptoms favor this diagnosis over some of the others, and this would be my first choice. Furthermore, by the law of averages I favor a malignant tumor. In my experience most of these tumors have been left-sided; so I would guess a left-sided tumor.

The second possibility I would consider is hyperplasia of the reticular cells of the adrenal cortex. This is a very interesting condition. It occurs mostly in females, and there is often a family history in the siblings. The typical story in females is usually as follows: together with certain characteristic changes in the external genitalia, the child is born with a large clitoris; she grows normally for the first one to eight years; she then develops axillary and pubic hair and grows rapidly, which suggests that androgen is being produced in excess somewhere in the body; finally, because of early epiphyseal union secondary to the excess of androgen, she stops growing at an early age. The net result is that her final height is about normal.

The reticular cells of the adrenal cortex have an interesting life history as worked out by Dr. Sam S. Blackman, Jr. The fetal reticular zone is well developed and apparently functioning; in the first two weeks of postnatal life it retrogresses but leaves behind some cells that give origin to the postnatal reticular zone; these cells lie dormant until about puberty, when they again show signs of activity and produce reddish-brown granules.

The life history of the reticular zone in cases with hyperplasia of the reticular cells is modified. In the first place the fetal reticular zone is increased; in the second place the cells that give rise to the adult reticular zone lie dormant only one to eight years, rather than up to the usual time of puberty.

We are now in a position to interpret the clinical manifestations. The anatomic changes at birth are undoubtedly the result of increased androgen production by the hyperplastic fetal reticular zone; the failure of axillary and pubic hair to develop in the first few months is due to the latent period when the fetal reticular cells lie dormant; finally, the rapid growth of axillary and pubic hair and of the external genitalia, if it does occur, is evidence of the recrudescence of the reticular cells.

The reticular-cell hyperplasia not only is much less frequent in the male but also runs a somewhat different course. It is very apt to be associated with Addison's disease, because of the encroachment of the reticular cells on the other layers of the adrenal cortex.

Against the diagnosis of hyperplasia in this particular case are the failure of the genitalia to be enlarged at birth and the absence of any clinical or laboratory findings suggesting Addison's disease (for example, the serum electrolyte values and electrolyte values for sweat). On the other hand the onset of pubic hair and of rapid growth at six years of age is about the right time one would expect if this were the diagnosis. Furthermore, there must be cases in the male in which the activity of the fetal reticular zone is not sufficient to cause noticeable enlargement of the male genitalia. Finally, there must be male patients who do not have Addison's disease. Adrenocortical hyperplasia is my second choice.

I will now turn to possible disorders of the Leydig cells. First, a word about testicular function: the hypothalamus, by way of a neurohumeral pathway, stimulates the anterior pituitary body to release follicle stimulating hormone (FSH) and lutenizing hormone (L.H.) These hormones, in turn, stimulate the testes, FSH leading to tubular development and LH controlling production of testosterone by the Leydig cell.

The first possibility we might consider here as a cause of increase in androgen production is a tumor of the Leydig cells themselves. This is very rare. No such tumor was felt. In the only case I know of with this diagnosis, the 17-ketosteroid excretion was over 1000 mg. as compared with 6.6 mg. per 24 hours in this case. I think we can dismiss this diagnosis as possible but unlikely.

One might consider a tumor of the pituitary body giving off too much LH. Such a diagnosis would seem reasonable, but I never heard of a pituitary tumor that did this.


The next possibility to consider is a disturbance in the hypothalamus, releasing FSH and LH and stimulating all functions of the testes. One meets this condition in the so-called "pineal syndrome." It is initiated, not by the hormone of the pineal body, but by any tumor in the region of the pineal body. Against such a diag-

nosis in the case at hand is the failure of the testes to be enlarged. The size of the testes is dependent almost entirely on the size of the tubules, and in the pineal syndrome development of the tubular, as well as the Leydig cells, occurs. There are no findings to suggest an intracranial lesion, and I think we can dismiss this diagnosis.


Finally, one comes down to the possibility of a selective overproduction of LH without FSH. This might cause stimulation of Leydig cells without stimulation of tubules. I had never heard of such a syndrome until recently when Dr. Nathan Talbot mentioned it. I leave this for him to discuss. Such a diagnosis would seem to be compatible with the findings in this case, and I might put it down as a third choice.

It is now the vogue to divide adrenocortical hormones into three functional types, one for each layer of the gland. The innermost reticular layer is thought to produce the gluconeogenic, "sugar" or "S" hormone; the outermost glomerulosa layer is thought to produce the salt and water, sodium or "Na" hormone. It may be of interest to see what evidence there is for overproduction of each one of these hormones by the tumor that I have diagnosed. We have already discussed the androgenic hormone. Against overproduction of the gluconeogenic hormone are the failure of the eosinophils to be decreased, the normal value for 11-17 corticosteroids and the rapid growth. In favor of some overproduction of the "na" hormone is the hypertension and the low chloride and high potassium values in the sweat. It would seem, therefore, that this tumor was producing a large amount of androgenic hormone, a normal amount of the gluconeogenic hormone and a slightly increased amount of the "Na" hormone.

Dr. Nathan B. Talbot: Our preoperative diagnosis was the same as that just given by Dr. Albright. In the differential diagnosis the condition of greatest interest was a syndrome that we have now seen in a number of young boys. It is characterized by precocious masculinization, starting in early life. The testes are of normal size for age. The urinary 17-ketosteroid output is either normal or slightly elevated, and the urinary gonadotropin (FSH) assay is apt to be negative for 3 and 6 mouse units per 24 hours. Exploration of the adrenal glands of such patients fails to reveal either a tumor or bilateral hyperplasia. On the other hand, testicular



Central origin of the vagus nerves (parasympathetic)



Abdominal autonomic plexus (sympathetic)

Control of Gastric Motility and Spasticity in Peptic Ulcer with Banthine®

"The need¹ for suppressing gastric motility and spastic states is . . . fundamental in peptic ulcer therapy. Since the cholinergic nerves are motor and secretory to the stomach and motor to the intestines, agents capable of blocking cholinergic nerve stimulation are frequently used to lessen motor activity and hypermotility."

Banthine² "has dual effectiveness; it inhibits acetylcholine liberated at the postganglionic parasympathetic nerve endings and it blocks acetylcholine transmission through autonomic ganglia."

It has been shown¹ to diminish gastric motility and secretion significantly as well as intestinal and colonic motility.

The usual schedule of administration in peptic ulcer is 50 to 100 mg. every six hours, day and night, with subsequent adjustment to the patient's needs and tolerance. After the ulcer is healed, maintenance therapy, approximately half of the therapeutic dosage, should be continued for reasonable assurance of nonrecurrence.

Banthine® (brand of methantheline bromide) is supplied in: Banthine ampuls, 50 mg.—Banthine tablets, 50 mg.

It is accepted by the Council on Pharmacy and Chemistry of the American Medical Association.

1. Zupko, A. G.: Pharmacology and the General Practitioner, GP 7:55 (March) 1953.

2. McHardy, G. G., and Others: Clinical Evaluation of Methantheline (Banthine) Bromide in Gastroenterology, J.A.M.A. 147:1620 (Dec. 22) 1951.

SEARLE Research in the Service of Medicine

biopsy reveals marked development of the interstitial cells of Leydig without more than minor evidences of testicular tubular development. It is believed that the precocity is due to a production of androgenic hormones by the Leydig cells and that these in turn have been precociously developed and activated by pituitary interstitial-cell-stimulating hormone (ICSH). It is further believed that the pituitary body has been activated to ICSH production by some sort of hypothalamic disturbance. Since most of these patients have shown no neurologic evidences of an intracranial lesion, it seems probable that the hypothalamic disturbance is of a functional nature rather than an organic, structural defect. Parenthetically, it may be added that the testicular interstitial cells occupy but a small volume; hence, it is possible for extensive interstitial-cell hyperplasia to occur without a clinically definite increase in gross testicular size. It is growth of spermatic tubules that accounts in the main for the increase in testicular size during adolescence.

So far as testicular size and masculine development were concerned, the patient under consideration corresponded to patients with the condition just described. On the other hand, he differed in the age at onset of precocity, which in his case was much too late for the above syndrome. He also differed in water and electrolyte status, for he had signs (questionable edema of the face, slight hypertension and cardiac enlargement) suggestive of a pathologic increase in the production of adrenocortical hormones of the type that cause sodium and water retention and potassium diuresis. To gain some objective information concerning the present patient's adrenocortical "Na-K" hormone status, Dr. William Locke applied a sweat electrolyte test on which he has been working recently. It might be interesting to hear the results of these studies.

Dr. William Locke: Methods for estimating the rate of secretion of the "Na-K" hormone are not so satisfactory as the methods for the other cortical hormones, as Dr. Albright and Dr. Talbot have said. In this case the hypertension may have indicated some overproduction of that hormone. The patient did not have hypochloremic alkalosis, which is sometimes associated with Cushing's syndrome and which has been attributed to an overproduction of the "Na-K" hormone in that condition. We have been studying variations in the electrolyte concentrations in

sweat, with a view to using them as an index of "Na-K" hormone production. This subject was first studied by Dr. J. W. Conn, who showed that desoxycorticosterone acetate has somewhat the same effect on sweat as it has on urine composition—that is, it reduces the concentration of chloride and sodium, and increases the concentration of potassium. In this case the sweat chloride was 10.0 milliequiv., and the potassium 13.2 milliequiv. per liter. Under the conditions of this test one would expect the chloride concentration in the sweat from a normal child to be perhaps three times what it was in this case and the potassium concentration to be approximately half what it was in this case. So we considered our findings to be suggestive evidence of an excessive production of the "Na-K" hormone.

CLINICAL DIAGNOSIS

Carcinoma of adrenal cortex.

DR. ALBRIGHT'S DIAGNOSIS

Carcinoma of adrenal cortex.

ANATOMICAL DIAGNOSIS

Carcinoma of adrenal cortex.

PATHOLOGICAL DISCUSSION

Dr. Oliver Cope: This patient had a tumor of the adrenal cortex. Why were the roentgenograms reported as showing no evidence of a tumor? The answer is that the tumor was small, not more than 3 cm. in its greatest diameter and scarcely large enough to cast a shadow. The roentgenograms were excellently clear, and a shadow was present on either side just above the kidney consistent with that case by the adrenal gland. The shadow on the right was denser than that on the left. In making the incision we anticipated that if a tumor were present it would be on the right side. This proved to be the case.

It is a curious thing, as Dr. Albright has pointed out, that the overwhelming majority of the hyperfunctioning adrenocortical tumors we have encountered have been on the left side. Left-sidedness to me makes no sense so I am always equally suspicious of the right. This is the second right-sided tumor of our series. Left-sidedness is presumably a statistical fluke and I suspect that with time and increased experience the left-sided predominance will also disappear.

Careful search was made at operation for evidence of malignant spread of the tumor. Grossly, the capsule of the tumor was unbroken. No

invaded lymph nodes were encountered. The tumor was excised in toto.

The tumor was transected after removal at the operating table. On the gross examination I made a diagnosis of cancer because of the unevenness of consistence in one area and distortion of the stroma. These gross findings have been present in other malignant adrenocortical tumors and consistently absent in the small, apparently benign, tumors that have been resected from patients with Cushing's disease. My experience is too limited for me to be certain but I have a hunch that in such tumors the gross pathology is more reliable than the microscopical. I am therefore fearful of recurrence in spite of my negative search for lymph-node metastases.

Dr. Tracy B. Mallory: The tumor was a well encapsulated nodule of quite variable color with many bright-orange-yellow spots in it, indicating that in portions of the tumor considerable lipoids were present. The microscopical section showed a very variable picture. Cells could be found that corresponded to all the various layers of the adrenal cortex. There were numerous lipoid-filled cells corresponding to the fascicular layer, and also many of the big rather homogeneous eosinophil cells that are found in the reticular layer. There was not a great deal of pigment in the cells in this case. The question comes up whether this tumor was benign or malignant. Encapsulation would be in favor of its being benign. In some parts of the tumor very atypical cells with giant multiple nuclei were found, which anywhere else in the body would certainly be evidence of a highly malignant tumor. Our experience in adrenal tumors has indicated that marked cellular atypicality of this type may be present in tumors that run a benign course. I think it very probable that if these slides were shown to a group of pathologists they would split fifty-fifty as to whether it was benign or malignant. My impression is that the outlook is relatively favorable although we did call it carcinoma. That may perfectly well be an error.

DIFFERENTIAL DIAGNOSIS IN THE RHEUMATIC DISEASES

From the number of papers appearing in the current literature, this is still a bothersome question, not yet entirely solved by the research departments of the pharmaceutical houses, though some of their literature would almost lead us to

think so.

Charley J. Smith, M.D., Denver, Colo., has a very excellent paper in the Rocky Mountain Medical Journal of January, 1953, confined to the differential diagnosis, the only comment on treatment being that "with the discovery and introduction into medical practice of Cortisone and ACTH, which are profoundly beneficial in a variety of rheumatic diseases, it has become more important than ever to establish an exact diagnosis in the early stages of these diseases . . ."

He discusses rheumatoid arthritis and rheumatic fever, degenerative vs. rheumatoid arthritis (with a diagnostic table), special types of rheumatoid arthritis, infectious arthritis, Reiter's syndrome, acute gout, traumatic arthritis. W.W.W.

INTERNATIONAL ACADEMY OF PROCTOLOGY

The International Academy of Proctology announces the establishment and award of a one year Proctologic Research Fellowship in the amount of \$1200.00. This Research Fellowship grant has been awarded to the Jersey City Medical Center, New Jersey, to be administered under the direction of Dr. Earl J. Halligan, Surgical Director of the Medical Center.

Dr. Halligan is a former International President of the Academy. The Board of Trustees of the International Academy of Proctology will vote another Fellowship grant of a similar amount at the time of the next Annual Meeting of the Academy. Thus, there will be at least two Research Fellowship studies in progress, in different institutions, under the auspices of the International Academy of Proctology.

WANTED

To associate with or to assist G.P. or Internist in the Area of Phoenix or Tucson. I am interested in internal medicine especially cardiology. I am an American by birth, of the Jewish faith, and am 46 years of age. I have an Arizona Medical license. I am a graduate of State University of Iowa year 1931; have been in active practice since 1932. I have had hospital in-patient and out-patient experience plus much post graduate work.

Write % Arizona Medicine Journal
424 Heard Bldg., Phoenix, Arizona

MEETING NOTICE

1953 Annual Convention of the National Society for Crippled Children and Adults, Nov. 12-14, Chicago.

The PRESIDENT'S *Page*

Through a series of coincidences I gained knowledge this summer of the existence of Spears Chiropractic hospital in Denver, and predict that you will hear more about it. A Tucson physician sent me a many-paged advertisement given him by a patient, made up in newspaper form of testimonials telling of "cures" where other medical and surgical skill had failed. Cancer, polio, multiple sclerosis and numerous other scourges are "cured or remarkably improved". While in Denver one newspaper reported that the wife of a Texas utilities man suffering from advanced cancer was transferred by special train from an eastern medical school to Spears and on advice of the attending chiropractor the special coach and engine made the last part of the trip from Chicago alone because she was in much pain.

I drove around the Spears grounds. It is already a big outfit and additional units are being built in series, probably to accommodate as many as 1,000 patients. Testimonials indicate that patients are being referred by chiropractors from all sections of the country. Tax litigation is in progress, the city maintaining the institution is being operated for profit.

The joker in this is that the physical set-up at first glance might pass for a VA hospital. This thought may have crossed the minds of the builders and the nation's chiropractors. The Denver papers during late August carried a few stories on an inside page about hearings being conducted by Senator Langer (Rep.) of North Dakota, chairman of a senate investigating sub-committee, who opened his Denver phase of committee investigation into an alleged "medical monopoly" and what he called the "conspiracy on the part of the press and the medical profession to discredit the practice of chiropractic." Senator Langer demanded that the VA provide ex-servicemen with chiropractic treatment at tax-payer's expense, denounced the A. M. A. for its stand against indiscriminate hospitalization for veterans with non-service connected disabilities, said he favored expansion rather than reduction of VA services by inclusion of chiropractic, and blasted the press for not giving his remarks front page space equal to that of Dr. McCormick's address before the Rocky Mountain Radiological Society. The Honorable Mr. Langer was a patient "incognito" for eleven days prior to the hearings, and according to Dr. Leo Spears "improved considerably." Testimony was taken from chiropractic "benefited" veterans suffering from "polio, multiple sclerosis and nerve disorders" by Mr. Langer, the lone member of the committee present at the hearings.

Edward M. Hayden, M.D.
President

Editorial

ARIZONA MEDICINE

Journal of

ARIZONA MEDICAL ASSOCIATION, INC.

VOL. 10 OCTOBER, 1953 NO. 10

EDITORIAL BOARD

R. Lee Foster, M.D. Editor-in-Chief, Phoenix
Harold W. Kohl, M.D. Assistant Editor, Tucson

ASSOCIATE EDITORS

William H. Cleveland, M.D. Phoenix
Howard D. Cogswell, M.D. Tucson
Louis G. Jekel, M.D. Phoenix
Frank J. Milloy, M.D. Phoenix
William H. Oatway, Jr., M.D. Tucson
Leslie B. Smith, M.D. Phoenix
Arie C. Van Ravenswaay, M.D. Tucson
W. Warner Watkins, M.D. Phoenix

COMMITTEE ON PUBLISHING

R. Lee Foster, M.D., Chairman Phoenix
Carroll C. Creighton, M.D. Flagstaff
Donald E. Nelson, M.D. Safford
Darwin W. Neubauer, M.D. Tucson

Robert Carpenter, Executive Secretary,
Arizona Medical Association, Inc.

ADVERTISING AND SUBSCRIPTION OFFICERS

J. N. McMEekin, Publisher and Business Manager,
424 Heard Building, Phoenix, Arizona

Eastern Representative
A. J. JACKSON, Director
State Journal Advertising Bureau
535 N. Dearborn St., Chicago 10, Illinois

CONTRIBUTORS

The Editor sincerely solicits contributions of scientific articles for publication in ARIZONA MEDICINE. All such contributions are greatly appreciated. All will be given equal consideration.

Certain general rules must be followed, however, and the Editor therefore respectfully submits the following suggestions to authors and contributors:

1. Follow the general rules of good English, especially with regard to construction, diction, spelling, and punctuation.

2. Be guided by the general rules of medical writing as followed by the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION. (See MEDICAL WRITING by Morris Fishbein.)

3. Be brief, even while being thorough and complete. Avoid unnecessary words. Try to limit the article to 1500 words.

4. Read and re-read the manuscript several times to correct it, especially for spelling and punctuation.

5. Submit manuscript typewritten and double-spaced.

6. Articles for publication should have been read before a controversial body, e.g., a hospital staff meeting, or a county medical society meeting.

The Editor is always ready, willing, and happy to help in any way possible.

HOSPITAL BEDS

The magnificent new St. Joseph's Hospital adds more than 100 hospital beds to the total number in Phoenix. Good Samaritan Hospital is in the process of adding more. If the old St. Joseph's Hospital is used in part a group

of more than 100 beds will be added.

But still there may be a shortage, a shortage which surely will increase as the population grows — as it certainly will.

The physicians can make a real contribution to the solution of this problem in two ways:

1. Do not enter patients into hospitals without sufficient reason.
2. Dismiss patients from the hospital at the earliest possible safe and satisfactory time.

There can be no doubt that at times the physician is put under considerable pressure to hospitalize patients who could do as well at home. This is especially true when patients have hospitalization insurance. But it is the responsibility of the physician to decide the need for hospitalization and not that of the patient nor the patient's family. And, in general, it is best to keep the patient at home during any illness which will respond equally well to home treatment.

It is hardly questionable that many patients are allowed to remain in the hospital for one or a few or even many days after they could properly be sent home. This is usually the doctor's fault, and may at times be due to sheer carelessness. Many hospital bed days could be saved by prompt dismissal of patients.

As pointed out above, the physician's problem in this regard is not always an easy one. If, however, he will keep constantly in mind the goal of using no more hospital bed days than necessary, he will perform a notable public service.

NOTICE

**ALL CONTRIBUTORS OF
ARIZONA MEDICINE SHOULD
HAVE THEIR MATERIAL IN THE
JOURNAL OFFICE NOT LATER
THAN THE 10th OF THE MONTH
PRIOR TO PUBLICATION IN
ORDER TO HAVE ARIZONA
MEDICINE REACH ITS READERS
ON OR BEFORE THE 10th OF
THE MONTH**

Material arriving after that date will be published the following month.

Arizona Pharmaceutical Page

THE SECOND CENTURY - 1852-1953

AS WE GLANCE BACK OVER THE PAST CENTURY OF OFFICIAL OPERATION OF PHARMACY IN THIS COUNTRY, THROUGH ITS PARENT ORGANIZATION, "THE AMERICAN PHARMACEUTICAL ASSOCIATION," WE ARE AMAZED AT THE TREMENDOUS ADVANCES OUR PROFESSION HAS BEEN PRIVILEGED TO PARTICIPATE IN. AS THE YEARS HAVE PASSED DURING THIS CENTURY OF PROGRESS WE HAVE SEEN UNTOLD CHANGES IN OUR SOCIAL AND ECONOMIC ORDER. THIS PRESENT GENERATION HAS LIVED AND IS LIVING THROUGH SOME OF THE MOST MOMENTOUS OF THESE CHANGES.

JUST HOW WE HAVE PROGRESSED IN THE 100 YEARS IS EVIDENT ALL ABOUT US. THE SINGLE PHARMACIST AND HIS APPRENTICE WORKING IN THE SHOP OF 1852 NOT ONLY AS COMPOUNDERS OF PRESCRIPTIONS BUT ALSO AS SMALL-SCALE MANUFACTURERS OF THE PREPARATIONS PRESCRIBED HAVE BEEN REPLACED BY A VERITABLE ARMY OF SPECIALISTS IN PHARMACY, PHARMACEUTICAL CHEMISTRY, PHARMACOLOGY, BACTERIOLOGY AND THE MANY OTHER CONTRIBUTORY ARTS AND SCIENCES. THESE NEW PRACTITIONERS HAVE RECEIVED THE MOST THOROUGH TRAINING IN THEIR PROFESSION. IN THE STATE OF ARIZONA A PERSON STARTING IN THE PROFESSION OF PHARMACY MUST SPEND A MINIMUM OF SIX YEARS IN THE STUDY OF IT BEFORE BEING ELIGIBLE TO BECOME LICENSED TO PRACTICE.

HIS COLLEGE REQUISITES INCLUDE ONE YEAR OF PRE-PHARMACY AND FOUR YEARS OF THE PROFESSIONAL COURSE LEADING TO THE BACHELORS DEGREE. FOLLOWING THIS THE APPLICANT IS REQUIRED TO COMPLETE ONE YEAR OF INTERNSHIP, UNDER THE SUPERVISION OF A PRECEPTOR PHARMACIST.

ORIGINALLY PHARMACY AND MEDICINE HAD A COMMON ORIGIN, GRADUALLY SPLITTING INTO TWO WELL DEFINED PROFESSIONS. AS SCIENTIFIC KNOWLEDGE DEVELOPED, IT BECAME INCREASINGLY DIFFICULT FOR ANY SINGLE GROUP TO KEEP PACE WITH PROGRESS IN ALL PHASES OF THE HEALTH FIELD. TODAY WE RELY ON THE PHARMACIST TO DEVELOP AND PREPARE THE MEDICINES REQUIRED TO MEET THE NEED SHOWN BY THE PHYSICIAN'S DIAGNOSIS.

BUT ALL OF THIS IS PAST HISTORY - WHAT OF THE FUTURE? IF WE ARE TO JUDGE THE COMING CENTURY IN THE LIGHT OF THE PROGRESS MADE DURING THE PAST CENTURY WE MUST HAVE A VIVID IMAGINATION AND BE ABLE TO VISUALIZE UNHEARD OF ACCOMPLISHMENTS. EIGHT YEARS AGO WE WERE PERMITTED TO DISPENSE PENICILLIN; THREE YEARS AGO WE RECEIVED OUR PORTION OF STREPTOMYCIN; EIGHTEEN YEARS AGO WE RECEIVED A STOCK OF SULANILIMIDE; TWO YEARS AGO CAME ACTH AND CORTISONE. WE COULD GO ON AND ON WITH A LISTING OF THE HORMONES THE ANTIHISTAMINES THE NEWER ANTIBIOTICS ETC. NEEDLESS TO SAY THE DISCOVERIES AND ADVANCES MADE WITHIN THE PAST TEN YEARS HAVE OVERSHADOWED ALL THOSE MADE IN THE PAST. AS WE CONTEMPLATE THE ADVANCES AND DISCOVERIES WHICH WILL BE MADE IN THE COMING CENTURY OUR IMAGINATION IS REALLY STAGGERED. YOUR DOCTORS WILL HAVE AVAILABLE WITHIN A VERY SHORT TIME MEDICATIONS TO SUCCESSFULLY TREAT MOST OF THE PRESENTLY KNOWN DISEASES. YOU MAY REST ASSURED YOUR PHARMACIST WILL HAVE KEPT ABREAST OF THE ADVANCES AND WILL BE READY TO SERVE YOUR NEEDS AS THE TIME ARISES.




They'd crowd an outdoor cafe...

all the patients who represent

the 44 uses for short-acting NEMBUTAL®

44 PATIENTS? Just look in the picture above. You'll find them all. And with every NEMBUTAL patient, with every NEMBUTAL use, these are the facts that you'll find the same:



1. Short-acting NEMBUTAL (Pentobarbital, Abbott) can produce any desired degree of cerebral depression—from mild sedation to deep hypnosis.
2. The dosage required is small—only about one-half that of many other barbiturates.
3. There's less drug to be inactivated, shorter duration of effect, wide margin of safety and little tendency toward morning-after hangover.
4. In equal oral doses, no other barbiturate combines quicker, briefer, more profound effect.

All are sound enough reasons for your prescription to call for short-acting NEMBUTAL. How many uses have you tried?

Abbott

1-512A

WHEN WRITING ADVERTISERS PLEASE MENTION THIS JOURNAL

TOPICS OF *Current Medical* INTEREST

RX., DX., AND DRS.

By GUILLERMO OSLER, M. D.

Dr. Arthur Kemmerer, professor of nutrition at the U. of A. College of Agriculture, gave a talk before the American Dietetic Ass'n. in Los Angeles. He is a member of the U. of A. arthritis research group. . . . His major comment concerning the use of ACTH and CORTISONE has not been generally publicized, and seems worth a mention here. Persons with ARTHRITIS apparently do not use PROTEINS in a normal manner. Use of the drugs enables them to metabolize protein, and the symptoms subside during that period. . . . On this subject, mas des pues.

The recently publicized 'DAILY DOUBLE' treatment for TB is double talk. . . . It came from Denver in September and is much the same as if a new, jazzy name for aspirin had been invented. The treatment (isoniazid, etc., plus gradual rehabilitation) is what most everyone else is using. The Denver people eliminate excessive bedrest, but continue therapy for 16 to 18 months. . . . The authors are Dressler and Middlebrook, and they must have a good publicity agent. It may be only coincidence but the name of the lab at the National Jewish Hospital is the 'Hearst Research Laboratories'. . . . Believe it or not, this is the same outfit which stirred up the hopeful last winter by saying that they were decreasing rest to as little as 1 to 3 months. Sanatoria would become obsolete. They back-tracked when investigated by the N T A. . . . This barefoot country Osler holds that such conduct ain't fair.

A further test of antibiotics for INTestinal AMEBIASIS has been made at Louisiana State by McHardy and Frye. . . . Terramycin leads the pack (91.5% success in 435 cases). Fumagillin, with its narrow spectrum and direct action is next (86% of 119 cases). Aureomycin cleared 83.4% of 697 cases. . . . It should be emphasized that extra-intestinal lesions (e. g., liver abscess) are not affected by the AB drugs.

Our old friend, the Florida Citrus Commission, sends out a flash bulletin about teen-agers. Their diets are deficient in VITAMIN C, from the rock-bound coast of Maine (Dr. Mary Clayton) to the sunny slopes of Texas (Dr. Pauline Mack). . . . The Commission has its own solution of course, but how about an Arizona angle? Send all the kids to school in Arizona and we will fill them up (6 to 8 ounces every morning, same like Florida) with Arizona orange juice. . . . The gums need Vit. C, they say, and it helps make the cells (as they say in Milwaukee) "tagedder schticken".

HEMOPHILIA is rarely seen in general practice. Surgery and other bloody procedures are usually feared and avoided. A recent description of bloodless DENTAL EXTRACTION is of some interest, and you'd hardly ever guess how it is done. . . . A rubber band is fitted tightly around the tooth. It gradually works its way down below the gingival margin and toward the end of the root. The tooth loosens and falls out, after periods ranging from 4 to 200 days. . . . No mention was made of a pain factor.

DR. ALVAREZ SAYS.—That he finds the evidence contradictory as to why some persons are long lived. . . . The Dunbars reported at St. Louis that longevity does not seem to be inherited. They studied the 300 cases of the 1500 White Americans who had lived over 100 years. . . . Dr. A. quotes the late Dr. Raymond Pearl, a great authority on the subject, who believed that if a man could choose his ancestors he could be sure of a long life. . . . Homologous twins often live lives which are nearly the same length. They are like 'duplicate clocks, wound to the same degree of tightness, which run down and stop at the same time'.

CYCLONES of the recent summer gave several HOSPITALS an acid test. They stood up well and learned a lot, some of it fast. . . . Providence Hospital in Waco, Texas, has 250 beds. They were presented with 215 cyclone victims in 2 hours. They admitted 115 and treated 100 as OPDs. Fifty of the latter were turned over to Connall Air Base hospital thru some fine civilian-military cooperation. . . . St. Joseph's Hospital of Flint, Michigan, also has 250 beds, but had a pre-cyclone census of 262. They gave first aid to 120 patients, admitted 78, and took 110 x-rays in one night. The success of the Flint operation was obtained by alerting the employees, gathering all of the staff, and calling in as many ex-employees as possible.

Ravits of St. Paul discusses 'DERMO-FUNGOUS ALLERGY' in MINNESOTA MEDICINE. He agrees with the work of Bloch and Jadassohn, who explain the sequence which can follow a trichophyte infection. . . . A person has a fungus infection (athlete's foot); he recovers; a generalized allergy develops to its height a few weeks after healing; it gradually declines thereafter; a person may react to the fungus or its products by skin reactions ('ids'). The lesions may be less violent and more eczematoid. . . . Skin-testing with an allergen (trichopytin) may be helpful in



THE HOSPITAL BENEFIT

Bulletin

Special

Published Bi-Monthly by the Hospital Benefit Association, First Street at Willetta, Phoenix

October, 1953

Question Quiz

Do you know the answers?

Q. Can doctors help expedite processing of claims?

A. Yes. In fact, the Association wants to pay claims quickly. It will do so if doctors will fill out and return promptly the claim form sent them by the Association.

Q. What standard does the Association use in paying fees?

A. The Association bases its schedule of benefits on the schedule which is used by the Arizona Industrial Commission and which was approved by a committee representing the Arizona Medical Society.

Q. Does the Association pay surgical fees to osteopaths?

A. No. Member benefits are paid only to licensed physicians (M.D.'s).

Q. May HBA members choose their own hospitals and doctors?

A. Yes. Member benefits are paid to any licensed hospital or physician (M.D.) anywhere in the world.

Q. Does the Association make an attempt to set the surgical fee?

A. No. This important decision is left entirely to the doctor and his patient. The actual cost is paid by the Association up to the amount shown in the schedule of fees.

Q. How does the doctor charge the Association for an X-ray?

A. He merely lists the charge for the X-ray on the claim form along with his own charges.

Q. Why does the Hospital Benefit Association have such a low rate of questionable claims?

A. The low rate is due to the fact that enrollment is handled properly by well-trained personnel. Members know just what benefits to which they are entitled.

HBA Surgical Plan Pays For Accident Treatment In Physicians' Offices

Doctors have been generous in their praise of the Hospital Benefit Association's Surgical Plan which contains many broad, generous benefits not offered in most other plans. They especially seem to like the fact that the Association pays for emergency treatment of accidents whether it is in the doctor's office, at the scene of the accident or in the patient-member's home. Of course, such treatments must be made within 24 hours of the accident. If X-rays for fractures, dislocations or for locating foreign bodies are required, as the result of an accident, the Association will pay for one X-ray, regardless of whether the X-ray is made at the doctor's office or at an X-ray laboratory.

Emergency accident treatment is included in the HBA Surgical Plan because the Association realizes the importance of this provision to both doctor and patient.

Everyday accidents — the kind that are most likely to happen to anyone at anytime — often do not require hospitalization but can be treated at the doctor's office. By including this provision, the Association enables many people to go to the doctor for minor injuries which otherwise might be neglected until they become serious.

Generous Benefits

Surgical fees are paid for members by the Association for fractures, dislocations, wound repairs and even examination of a suspected injury as the result of an accident. The HBA Surgical Plan also includes benefits to cover such non-surgical accident treatment as removal of a foreign body from a patient's eye, insect stings, accidental poisoning, etc.

X-rays in the Doctor's Office

Fees paid for X-rays of accidental injuries, whether they are made in the doctor's office or at the laboratory, range from \$6.00 to \$45.00, depending on the type of X-ray. This schedule of fees is based on the schedule which is used by the Arizona Industrial Commission and which was approved by a committee representing the Arizona Medical Society.

Simple Form Saves Time

The claim form which the Association sends the doctor is very popular because of its simplicity. It eliminates the red tape often encountered in establishing claim data. The doctor merely fills out the short form, listing his diagnosis, treatment, charges and answers a few simple questions.

Doctors Commend HBA TV Newsreel

The Association is pleased to report it has received many favorable comments from doctors on its TV show, the HBA Newsreel. This half-hour program, featuring noteworthy current news events, is telecast every Monday at 8:30 P.M. on KTYL-TV, Channel 12.



"Let him yelo — don't spoil him because he's an only child."

diagnosis, but is rarely of use in desensitizing. Treatment must usually consist of dull local applications.

It seemed a good idea to write a comment on the newest DENTAL PROTECTIVE PROCEDURE. The Sat. Eve. Post had a wide-eyed article on the ANTI-ENZYME approach, described by its Chicago originators as fabulous and amazing, as well as a continuous barrier to decay. . . . This merry-go-round is the same one we rode on 3 years ago ('ammoniated') and about 2 years ago ('chlorophyll'), so we were all ready to look with a jaundiced eye, when crash boom! whee! out came full page newspaper ADVERTISEMENTS referring to the article in THREE magazines. They all had just happened to be about anti-enzymes. . . . The ad also just happened to be about Antizyme, a new dentifrice by Listerine. The Lamberts ride again! . . . So we grabbed the brass ring and bought a tube, not for protection or science, but just to see how it tastes. . . . Mint.

The English feel that isoniazid offers a special help in TUBERCULOSIS OF THE KIDNEYS (Dick, in The Lancet). It is given credit for producing 'patholytic' miracles which most Americans have not yet admitted. . . . Caseation is said to be absorbed, lesions vascularized, and fibrosis prevented in the surrounding area. This is a reversal of the usual TB process. . . . They stress that "none of the antituberculosis drugs should be used ALONE."

Another new anti-TB drug is still too young to be well studied. The material is called HES, for hydroxethyl sulfone. It was synthesized by the National Institutes of Health Laboratories, and studies by Payne and colleagues of Howard University in Washington, D. C. . . . The place of HES in therapy, or its relationship to other drugs, or its relative toxicity is not certain enough for us to trifle with. . . . Hold off. Wait and see.

..STREPTOMYCIN sometimes produces a violent allergic reaction. In theory the drug should have been given on a previous occasion, but the patient denies it, or can't remember it, or doesn't know about it. . . . The latter case may have occurred when a double-drug combination was given as a shotgun therapy for some sore of infection. . . . There is a lot of evidence against the use of two drugs, especially when it is needless or careless, and especially when they are called by some cryptic trade name. Use 'em simple, we say, unless absolutely indicated, and that includes SM.

Drs. Clarke, Zahn, and Holmes of Seattle believe they have proof that A GOOD SCARE may help TUBERCULOSIS to heal. . . . They have clinical evidence that fright has seemed to turn the tide in many cases. They intend to use the method for a trial, and to further correlate the

relations of fear and TB and the adrenal gland. . . . Chemical tests of adrenal activity have been somewhat correlated with mental states, but the main work remains to be done. . . . We expect to see Dr. Zahn in Utah soon, and will find out the progress.

Dr. Frank Meleney, the antibiotic man from New York, sometimes plays a guest-spot in this journal. . . . He has recently clarified the BOIL AND CARBUNCLE situation, especially for the non-surgeon who hesitates at direct, local therapy. . . . Penicillin or bacitracin may be injected into the CENTER of furuncles and carbuncles. It also can be placed in the periphery of a wide zone of induration which may be present around the latter, in combination with a local anaesthetic, just as has been done by pioneers in the past couple of years.

RESINS have been described which absorb various electrolytes from the intestinal tract. They now sell one ('Resion', by The National Drug Co.) which is supposed to DETOXIFY. . . . It is composed of polyamine methylene resin and two aluminum silicates. It is said to be effective in diarrhea, and in gastroenteritis, and in the nausea and vomiting of pregnancy. They use the term 'polyphasic absorbent'. . . . It sounds reasonable in this day and age, but data are not sufficient for comparison with the simple kaolin-pectates, etc., nor whether the stuff may pick up too many chemical items.

A recent story in this column about a Los Angeles and a San Francisco chest surgeon has smoked out another 'gay' story. It concerns Dr. B., a Los Angeles surgeon who also is well-known in Arizona. . . . Dr. B's family has two cats. The 'outside cat' was hit by a car, taken to a veterinarian's and recovered except for a diaphragmatic hernia. . . . The vet was so delighted at this finding that he kept the cat for days and weeks. Mrs. B. called repeatedly, and the cat was always about to have another x-ray. She was told that the cat needed an operation, to which she replied that her husband did that sort of thing. One remark led to another, and the impasse was broken only when Mrs. B. arranged to get the cat home by allowing the vet to 'scrub' for Dr. B. during the hernia repair!

Electrocardiographer, certified, desires to interpret EKG's by mail. Replies by return mail day of receipt. \$1.25 per interpretation.

Write %
ARIZONA MEDICINE JOURNAL
424 Heard Bldg., 112 N. Central
Phoenix, Arizona

Your Official Professional Group Accident and Sickness Plan

Approved and recommended by Council Of

THE ARIZONA MEDICAL ASSOCIATION, INC.

Provides Maximum Protection at Minimum Cost
World Wide Coverage

IT PAYS YOU:

\$300 a Month for Total Disability by Accident up to 5 years	\$2,500 Accidental Death	\$7.00 a Day for Hospital Plus \$25 for Miscellaneous Expenses
\$150 a Month for Partial Disability by Accident up to 6 months	\$10,000 Dismemberment and Loss of Sight	\$5.00 a Day for Graduate Nurse, at home
\$300 a Month for Sickness up to 2 years		

LOW SEMI-ANNUAL PREMIUMS

Through Age 49—\$49.80 Ages 50 through 59—\$56.60 Ages 60 to 65—\$70.05

NO AGE LIMIT FOR RENEWAL

Policy Cannot Be Terminated Except For

- | | |
|-----------------------------|--------------------------------------|
| 1. Non-payment of premium | 3. Loss of membership in Association |
| 2. Retirement from practice | 4. Termination of master policy |

For additional information and official application contact

SIMIS INSURANCE SERVICE AGENCY

State Representatives

NATIONAL CASUALTY COMPANY

DWIGHT McCLURE
Telephone ALpine 3-1185

GEORGE B. LITTLEFIELD

W. J. WINGAR
407 Luhrs Building, Phoenix

PAUL H. JONES INSURANCE AGENCY

Pima County Representative

617 N. Stone Avenue, Tucson, Arizona

Telephone: Tucson 2-2803

Our members are dairymen whose business is supplying good milk for

WEBSTER'S

milk and milk products.

We are aware of the importance of good milk to good health and of our obligation to supply a product which will merit your confidence.

ARIZONA MILK PRODUCERS
422 Heard Building Phone ALpine 3-0893

METAL OFFICE FURNITURE

G. F. STEEL DESKS

ALUMINUM CHAIRS

CARDINEER ROTARY & VICTOR

VISIBLE FILES

Heinze Bowen & Harrington, Inc.

228 West Washington St.

Phone ALpine 4-4179

PHOENIX ARIZONA

WHY BUY

WHEN WE SUPPLY

**Everything in linens and
uniforms for the physician
and his staff**

CITY LINEN SUPPLY INC.

333 N. 7th Ave. — Phoenix, Arizona

ALpine 3-5175

Woman's AUXILIARY

THE PLEDGE OF LOYALTY TO THE WOMAN'S AUXILIARY

I PLEDGE MY LOYALTY AND DEVOTION TO THE WOMAN'S AUXILIARY TO THE AMERICAN MEDICAL ASSOCIATION. I WILL SUPPORT ITS ACTIVITIES, PROTECT ITS REPUTATION AND EVER SUSTAIN ITS HIGH IDEALS.

This pledge was adopted at the annual convention in Chicago in June 1943.

What is loyalty? There is probably no word in the English language more widely used. But do we appreciate its full meaning? Being true to one's inmost self and acting accordingly is loyalty. Webster says it is "Fidelity to a superior or to duty, allegiance, homage." The loyalty and allegiance of the Woman's Auxiliary is to the medical profession.

From the time of its organization in 1922, the membership of the Auxiliary has steadily increased, and the importance and scope of its activities in the advancement of health have become broader and more varied. It is the function of the Woman's Auxiliary, under the direction of the American Medical Association, to aid in the promulgation of those principles which will improve public health. Also, it is important that it be realized that one of the chief objects of the Woman's Auxiliary, when it was organized, was to promote friendliness among physicians' families.

Today there is need, as never before, to adhere to and to promote the objectives for which we stand as an Auxiliary to the American Medical Association. We should assume our share of responsibility for safeguarding the ideals of American Medicine at all times and under all conditions.

As an Auxiliary we should look to our advisers for direction and remember that, as an organization, we have a oneness of purpose which is definite and impersonal, yet at the same time unobtrusive and regulated. Let us hold to our ideals as we pledge our unified effort, in a spirit of loyalty and devotion, to the members of the great humanitarian profession which we have the privileges of association.

Mrs. David W. Thomas, Member of

the Publications Committee, and Past President, Woman's Auxiliary. (Taken from the August issue of the *Bulletin* of the Woman's Auxiliary to the American Medical Association.)

LEGISLATION REPORT

The Woman's Auxiliary to the American Medical Association has been called upon to assist with three major pieces of legislation during the coming year. These are: 1. The Bricker Resolution; 2. Bills allowing the deduction from income of expenses for post-graduate courses and advanced professional training, similar to H.R. 4393 by Davis of Georgia; 3. Reed-Keogh type bills, allowing tax deferment for retirement funds for the self-employed.

Senator Bricker desires a constitutional amendment which would limit the treaty powers of our government as they are at present set up, and to "protect the United States Constitution and the American system of government established in the Constitution, including the Bill of Right, against subversion by the Treaty Power; that will prevent treaties from becoming internal law within the United States, except to the extent made so by congress, just as in the case of any other federal statute.

The Woman's Auxiliary has been informed that the International Labor Organization has approved certain treaties or covenants which are in opposition to the accepted custom in the United States. These covenants are to be presented to the Congress of the United States for ratification. These covenants are the motive behind Senator Bricker's proposed amendment.

At the 30th Annual Meeting of the Woman's Auxiliary to the American Medical Association in June of this year 1953, it was resolved by the Woman's Auxiliary that they go on record as disapproving such treaties and covenants and that they favor the proposed amendments to the Constitution of the United States relating to the ratification of treaties and covenants.

There are erudite proes and cons regarding this resolution but the present amended bill (Please turn to Page 382)

CHIROPODY, A HAND MAIDEN OF MEDICINE. Did you know this is what the Judicial Council of the American Medical Association says about chiropody? In 1939 this Council ruled that the practice of chiropody is not a cult practice as some have believed but "is rather a practice ancillary — a hand maiden — to medical practice in a limited field considered not important enough for a doctor of medicine to attend and therefore too often neglected. General opinion seems to be that chiropody fairly well satisfies a gap in medical care that the profession has failed to fill." Several large hospitals, as well as the Mayo Clinic, have chiropodists on their staffs, and the Mayo Clinic thinks so well of its chiropodist (Tarara) that a dinner was held in his honor on the occasion of his appointment as a voting member of the Clinic staff.

Read more about chiropody and chiropodists in the Delaware State Med. Journ., June, 1953, article by Friedman on "Dermatology and the Chiropodist," and the article by Sindoni on "The Role of the Chiropodist in Diabetes."

W.W.W.

FOR RENT

Two Doctors, can have ideal location on the East side near McDowell Road.
At Very reasonable figures. Good Parking.

See or Call

Robert Rae, Real Estate Broker
1414 E. McDowell Rd.
Telephone AL 4-1636 — Phoenix

Monty Lebert's Nursery And Flower Shop

Artistic Flower Arrangements for Every Occasion

Assorted Potted Plants
Dish Gardens
Beautiful Ceramics

We Specialize In Planter Box Materials

531 W. Thomas Road Phone AM 6-2401
Phoenix, Arizona

In very special cases
A very superior Brandy



Specify

84 PROOF

HENNESSY

THE WORLD'S PREFERRED
COGNAC BRANDY

For a beautifully illustrated book
on the story of Hennessy, write—

Schieffelin & Co., Dept. HT, 30 Cooper Square, N. Y. 54

Auralgan
In Acute Otitis Media
Rhinalgan
Safe Nasal Decongestant
O-tos-mo-san
In Fungus and Bacterial Ears

PAUL DESILETS

Representative

515 Ocean Ave. — Seal Beach, Calif.
DOHO CHEMICAL CORP.
100 Varick Street, New York 13, N. Y.
Telephone Worth 2-7707

TRUSSES — SURGICAL SUPPORTS

GROVE'S

Surgical Supports Store
3123 N. Central — Phone CR 4-5562
Lady and Man Fitter — Fitted Exactly As You Order
Across from Park Central Development On Central
Avenue — Private Fitting Rooms
Free Parking — Open Wednesday Nights

ARIZONA LIBRARY BINDING COMPANY

Binders of Medical Journals and Books
We Pay Freight Charges on Return Shipments
Prompt Service

Arnold A. Couturier

307 West Monroe — Phone ALpine 3-1861
Phoenix, Arizona

has been approved by the American Bar Association and the American Medical Association and because of this bill will have a rough road ahead it is urged that we exert continuous pressure so that the bill will be considered and favorably reported by the Senate. The same consideration should be given H. J. Res. 280, by Reed of Illinois which is identical with the Bricker Resolution.

On the subject of taxation there are two points of particular interest to the Medical profession. The first is in regard to the deduction of postgraduate education expenses. The second subject (Reed-Koegh) which is of particular importance is No. 36 on the agenda. This is the question of allowing tax deferral for the establishment of retirement funds for the self-employed. This legislation would allow the physician to set aside, on a tax deferred basis, up to ten per cent of his income, but not to exceed \$7,500.00 a year, in a trust fund, from which he may withdraw the money on retirement, paying taxes on it at that time. Mrs. Edgar E. Quale our National Legislation Chairman tells us that "all the old bills on

socialized medicine are back in the hopper, and we must put the fight against socialized medicine on our agenda as permanent unfinished business." Our obligation as dutiful American citizens will not permit us to relax our vigilance or let down our guard in our struggle to help establish and support legislation that will benefit the greatest number of our fellow citizens. Let us valiantly and consistently assist in maintaining the freedom which we now enjoy.

Mrs. Charles L. von Pohle, Chandler, Arizona, State Legislation Chairman, Delegate to the Annual Convention of the Women's Auxiliary to the American Medical Association.

RADIOLOGIST 29; desires association with established Radiologist; hospitals; group or clinic. Southwest.

Write:

Arizona Medicine Journal
424 Heard Bldg., 112 N. Central Ave.
Phoenix, Arizona

LABORATORIES

PROFESSIONAL X-RAY AND CLINICAL LABORATORY

507 Professional Bldg.
Phoenix, Arizona
Phone ALpine 3-4105

AND

MEDICAL CENTER X-RAY AND CLINICAL LABORATORY

1313 North 2nd Street
Phoenix, Arizona
Phone ALpine 8-3484

DIAGNOSTIC X-RAY

X-RAY THERAPY

RADIUM THERAPY

CLINICAL PATHOLOGY

TISSUE PATHOLOGY

ELECTROCARDIOGRAPHY

BASAL METABOLISM

R. Lee Foster, M.D., Director John W. Kennedy, M.D., Radiologist
W. Warner Watkins, M.D., Radiologist
Diplomates of American Board of Radiology
Lorel A. Stapley, M.D., Consultant Pathologist

LABORATORIES

Medical Arts Bldg.
543 E. McDowell Rd.

PHOENIX, ARIZONA

Telephone
ALpine 8-1601

The Diagnostic Laboratory

A Complete Analytical and Laboratory Service To The Medical Profession of Arizona

Protein Bound Iodine
Blood Cholinesterase
17-Ketosteroids
Corticosteroids
Phosphatases
Vitamin Determinations
Blood Volume
Blood pH Values
Electrolytes
Toxicology
Autopsies
Papanicolaou Stains
Liver Function Tests
Porphyrins

Streptolysin Titres
Rh Antibody Titres
Quantitative Serology
Heterophile Titres
Autogenous Vaccines
Hematology
Bacteriology
Parasitology
Gastric Analysis
Friedman Tests
Frog Pregnancy Tests
Mycology
Enzyme Chemistry
Spectroscopic Analysis

Radiography
Pelvimetry
Electrocardiography
Basal Metabolism
Vital Capacity
Salpingograms
Cholecystograms
Bronchograms
Gall Bladder Series
G. I. Series
Pyelograms
Myelograms
Cystograms
Electrometric Analysis

Maurice Rosenthal, M.D.
Diplomate, American
Board of Pathology

◆ Douglas D. Gain, M.D.
Diplomate, American
Board of Radiology

◆ Ernest H. Price, M.D.
Diplomate, American
Board of Radiology

◆ George Scharf, M.D.
Diplomate, American
Board of Pathology

G. O. HARTMAN, M.D.

PATHOLOGICAL LABORATORY

20 E. Ochoa St. Phone: 3-4861

TUCSON, ARIZONA

Professional X-ray and Clinical Laboratory

Successor To

PATHOLOGICAL LABORATORY

507 Professional Bldg.

Phoenix, Arizona

Phone ALpine 3-4105

DRS. FOSTER, WATKINS and KENNEDY

THIS SPACE FOR SALE
FOR INFORMATION AND RATES
write to

ARIZONA MEDICINE

424 Heard Bldg.

Phone ALpine 2-4884

PHOENIX, ARIZONA

MEDICAL CENTER X-RAY AND CLINICAL LABORATORY

1313 N. Second St.

Phoenix, Arizona

Phone ALpine 8-3484

DRS. FOSTER, WATKINS and KENNEDY

SANATORIUM DIRECTORY

LIVERMORE SANITARIUM



• The Hydropathic Department devoted to the treatment of general diseases, excluding surgical and acute infectious cases. Special attention given functional and organic nervous diseases. A well equipped clinical laboratory and modern X-ray Department are in use for diagnosis.

• The Cottage Department (for mental patients) has its own facilities for hydropathic and other treatments. It consists of small cottages with homelike surroundings, permitting the segregation of patients in accordance with the type of psychosis. Also bungalows for individual patients, offering the highest class of accommodations with privacy and comfort.

GENERAL FEATURES

1. Climatic advantages not excelled in United States. Beautiful grounds and attractive surrounding country.
2. Indoor and outdoor gymnastics under the charge of an athletic director. An excellent Occupational Department.
3. A resident medical staff. A large and well-trained nursing staff so that each patient is given careful individual attention.

Information and circulars upon request.

Address: O. B. JENSEN, M.D.
Superintendent and Medical Director
LIVERMORE, CALIFORNIA
Telephone 313

CITY OFFICES:

SAN FRANCISCO
450 Sutter Street
GARfield 1-5040

OAKLAND
1624 Franklin Street
GLencourt 1-5988

McKEE REST HOME

Aged & Convalescents — Home-like Atmosphere

24 Hour Nursing Care

Good Meals

644 E. 4th St. — Phone 4-1971 — Tucson, Ariz.

HILLCREST SANATORIUM

- Aged and Convalescents only.
- Cheerful Private Rooms.
- Reasonable Rates.
- 24 Hour Nursing Service.
- Non-Contagious — Non-Alcoholics — Non-Addicts

Phone 4-1562

No. 3rd Ave & Adams

Tucson, Arizona

HIGHLAND MANOR

(Mr. & Mrs. Marley Karns)

- Convalescent.
- Personalized Diets.
- 24 Hour Nursing Care.
- Located in a Quiet Zone.

1411 E. Highland Ave. Phoenix, Arizona
Telephone AM 5-2552

EVANS REST HOME

5255 N. 43rd Avenue, Glendale, Arizona
Telephone ALpine 4-6511 or YE 7-8335

Ethical — Efficient

24 hour care for YOUR patients
Any non-contagious case treated as you direct




Catalina

ARIZONA'S FINEST SANATORIUM
FOR NON-CONTAGIOUS DISORDERS

Open Medical Staff
Flexible Rates
Open all year
Your inspection invited

AMherst 6-7238
5055 N. 34th St.
Phoenix

Location: East Camelback Road and North 34th St.



SANATORIUM DIRECTORY—Cont'd)

ALCOHOLISM

There is, today, a rapidly increasing, nation-wide awareness of alcoholism as a disease, as a major health problem and as a responsibility of the medical profession.

Physicians throughout the country are recognizing the need of special facilities for the proper care and treatment of the victims of this disease.

In THE FRANKLIN Phoenix has a modern center for care and treatment of acute and chronic cases of alcoholism.

All treatment at THE FRANKLIN is under the supervision of a licensed physician. Patients may retain their own physicians or may be attended by a staff physician of THE FRANKLIN.

Nursing care is continuous, 24 hours a day, under supervision of a registered nurse; with a staff of understanding, experienced people.

Hospital License No. 71

The Franklin Hospital, Inc.

ALCOHOLIC PROBLEMS - EXCLUSIVELY

367 N. 21st Ave., Just Above Van Buren

Telephone ALpine 3-4751

Phoenix, Arizona

VALLEY OF THE SUN REST HOME

PHOENIX, ARIZONA

126 W. Maryland

AMherst 5-9055

Bed Patients — Convalescents
Accommodations for Aged and Confused

BUTLERS REST HOME

- Bed Patients and Chronics.
- Excellent Food.
- Television.

802 N. 7th St.

Phoenix, Arizona


Telephone AL 3-2592

LA SIESTA LODGE

A clean and comfortable Rest Home

1032 E. Camelback Rd. — Phoenix, Arizona

Telephone AM 6-7473



MEDICAL STAFF

CHARLES W. THOMPSON, M.D., F.A.C.P., Director

CLIFTON H. BRIGGS, M.D., F.A.C.S., Associate Director

ETHEL FANSON, M.D.

CARLOS F. SACASA, M.D.

DOUGLAS R. DODGE, M.D. HERBERT A. DUNCAN, M.D.

KENNETH P. NASH, M.D.

INTERNAL
MEDICINE
and
NERVOUS
DISORDERS

PASADENA, CALIFORNIA

SANATORIUM DIRECTORY

GLENDALE NURSING HOME

Arizona newest, modern nursing home.

- Convalescent
- Custodial
- 24 Hour Nursing Care
- Special Diets. Quiet.

Lat. 16 $\frac{3}{4}$ and Glendale Avenue

Phones: AMherst 6-7001 — YELlowstone 7-7064
Glendale, Arizona
(Ray and Ruth Eckel)

DESERT REST HOME

Convalescent Tubercular

- Reasonable Rates
- 24 Hour Nursing Care
- Excellent Care

409 E. Townley Ave. — Phone WINDsor 3-3689
Phoenix, Arizona (Sunnyslope)
(Ray and Ruth Eckel)

MESA SANITARIUM

Convalescent — Ambulant

Physician and Registered Nurses
On 24 Hour Duty

Allene Lauck, R. N.—Prop.

Telephone WO 4-3780

42 South Country Club Drive
Mesa, Arizona

THIS SPACE FOR SALE

FOR INFORMATION AND RATES
write to

ARIZONA MEDICINE

424 Heard Bldg.

Phone ALpine 2-4884

PHOENIX, ARIZONA

AMBULANCE DIRECTORY**A & A AMBULANCE CO.**

Tucson's Only Exclusive Ambulance Service

Oxygen Therapy

Day or Night

PHONE 3-3645

405 North 9th Ave.

Tucson, Arizona

MEDICAL SUPPLY DIRECTORY**ARIZONA MEDICAL SUPPLY CO.**

PHONE 3-7581

1025 E. Broadway — Tucson, Arizona

Martin O. Kerfoot

Geo. F. Dyer

MEDICAL BOOKS**ROBERTA DAVEY HALL**

MEDICAL BOOKS

Of All Publishers

PROMPT HANDLING OF ALL ORDERS

To order: Telephone AMherst 5-1062

40 East Rose Lane

Phoenix, Arizona

THIS SPACE FOR SALE

write to

ARIZONA MEDICINE

424 Heard Bldg.

Phone ALpine 2-4884

PHOENIX, ARIZONA

THIS SPACE FOR SALE
FOR INFORMATION AND RATES

write to

ARIZONA MEDICINE

424 Heard Bldg.

Phone ALpine 2-4884

PHOENIX, ARIZONA

THIS SPACE FOR SALE
FOR INFORMATION AND RATES

write to

ARIZONA MEDICINE

424 Heard Bldg.

Phone ALpine 2-4884

PHOENIX, ARIZONA

DRUGGISTS' DIRECTORY

WAYLAND'S**Wayland's Prescription Pharmacy****13 E. Monroe Street****Phone ALpine 4-4171****PHOENIX, ARIZONA****FREE DELIVERY**
MARTIN DRUG CO.
Rexall
28 Registered Pharmacists

Tucson

Casa Grande

PRESCRIPTION

Complete line of
 Hospital Beds, Crutches, Trusses and
 Surgical Garments

THE PRESCRIPTION SHOP

45 East Broadway

Phone 3-4701

TUCSON

D. F. Scheigert

L. J. McKenna

SOLANO PHARMACY

7th Ave. & Bethany Home Rd. — CR 4-2452

MELROSE PHARMACY
 4320 N. 7th Ave. AM 5-8411
 PHOENIX

THIS SPACE FOR SALE
 FOR INFORMATION AND RATES

write to

ARIZONA MEDICINE

424 Heard Bldg.
 Phone ALpine 2-4884
 PHOENIX, ARIZONA

MacALPINE
DRUG CO.
A complete line of . . .

PHOTOGRAPHIC SUPPLIES
 COSMETICS
 LIQUOR AND
 PRESCRIPTION DEPT.

Prompt FREE delivery

2303 North 7th St.,
 PHOENIX, ARIZONA

phones: ALpine 1-2606
 or ALpine 2-1573

* THE REXALL STORE

DRUGGISTS' DIRECTORY

Where Your Doctor Speaks and Your Druggist Serves

SIMON'S DRUGS

Prescriptions • Hospital Supplies • Sick Room
Supplies • Pharmaceuticals • Baby Needs
Trusses • Crutches • Abdominal Supports

2829 W. Van Buren—Phones AP 8-1611 - AP 8-2662

Phoenix, Arizona

Ample Parking Space — City-wide Free Delivery

LAIRD & DINES

The REXALL Store

Reliable Prescription Service

WOodland 7-2922

Mill Ave. & 5th

Tempe, Arizona

BEVERLY BURKE

PRESCRIPTION DRUG STORE

MEDICAL CENTER

1313 N. 2nd St. — Ph. ALpine 8-2706

Phoenix, Arizona



ALpine 3-2148

McCreary's
DRUG CO.
Central Ave. at McDowell

THIS SPACE FOR SALE
FOR INFORMATION AND RATES
write to

ARIZONA MEDICINE

424 Heard Bldg.
Phone ALpine 2-4884
PHOENIX, ARIZONA

CREIGHTON PHARMACY

Telephone BR 5-2441

2345 E. McDowell Rd. — Phoenix, Arizona

CHARLES E. BILL M. GERTRUDE BILL

JOHNSON'S DRUG STORE

PRESCRIPTIONS

"Service you will like"

Corner Speedway and Park Avenue
Phone 2-8865 Tucson, Arizona

ENSMINGER PHARMACY

RELIABLE PRESCRIPTIONS

121 North Cortez

Phone 188

Prescott, Arizona

FAIRMONT PHARMACY

Telephone BRidge 5-1331

3231 E. McDowell Rd.

Phoenix

PULLINS

Prescriptions

400 E. Glendale

Phone YE 7-9848

Glendale, Arizona

DRUGGISTS' DIRECTORY

EUREKA DRUGS

PRESCRIPTIONS

"The Friendly Drug Store"

R. L. Gibson, Prop.
Phone 2-7153 Tucson, Arizona**FELSHER PRESCRIPTION
PHARMACY**Sick Room Supplies — Vitamins
650 N. First Avenue — Phone ALpine 3-2070
Phoenix, Arizona**MEDICAL SQUARE PHARMACY**

PRESCRIPTION SERVICE

P. M. Corke Phone 5-3371
1616 No. Tucson Blvd. — Tucson, Arizona**EVERYBODY'S DRUG COMPANY**

Prescription Druggists

Phones: WO 4-4587 — WO 4-4588
Mesa, Arizona

Open 8 A.M. to 11 P.M. Daily & Sunday

Broadway Village Drug Store

PHONE 5-2631

Broadway at Country Club Road
(Free Delivery)

TUCSON ARIZONA

NATIONAL PHARMACY

Prescriptions

Phone 2-9779 340 Ajo Way
Tucson, Arizona**MONTE BLISS****Frontier Village Drug Store**

"Your Reliable Neighborhood Prescription Pharmacy"

1700 N. Maple Blvd. Phone 5-5252
Tucson, ArizonaTHIS SPACE FOR SALE
FOR INFORMATION AND RATES
write to**ARIZONA MEDICINE**424 Heard Bldg.
Phone ALpine 2-4884
PHOENIX, ARIZONATHIS SPACE FOR SALE
FOR INFORMATION AND RATES
write to**ARIZONA MEDICINE**424 Heard Bldg.
Phone ALpine 2-4884
PHOENIX, ARIZONA**STONE AND 3RD PHARMACY**

749 N. Stone — Phone 3-6041

ENCANTO PARK DRUG CO.3352 E. Speedway — Phone 5-3102
Tucson, Arizona**FLORES & SON**

(FARMACIA FLORES)

"Your Nyal Service Drug Store"
W. Congress and Meyer Sts. Phone 3-3362
Tucson, Arizona**MODERN RX PHARMACY**

TELEPHONE 20

NOGALES

ARIZONA

MINERAL WELLS DIRECTORY*Buckhorn*

- 27 Private Baths — 8 Whirlpool Baths
- 9 Massage Rooms
- 2 Lounge Rooms
- 10 Acres Beautiful Grounds

Open 8 a.m. to 8 p.m. Daily

Natural Hot Mineral Baths

(7 mi. East on Apache Trail)

Phone WOODland 4-7316 — Mesa, Arizona

X-RAY COMPANY**GENERAL ELECTRIC CO.****X-RAY DEPARTMENT****FOR SALE**Used G. E. 200 MA R&F Unit.
Hand Tilt Table, Good Condition.
General Electric Co. X-Ray Dept.

Ask for M. A. Schroeder — ALpine 4-0181

DOCTORS' DIRECTORY

DOCTORS DIRECTORY ESTABLISHED 1920

ALpine 3-4189

Emergency calls given special attention. We will locate your doctor before or after office hours.

BERTHA CASE, R.N., Director

ADA JOY CASE

1541 East Roosevelt
Phoenix, Arizona

DOCTORS CENTRAL DIRECTORY

Minnie C. Benson, R.N., Manager

For Emergencies or in Absence of Your Doctor

CALL 5-1551

At Your Service 24 Hours Daily

E. Hedrick Dr. Tucson, Arizona
"Established 1932"

NURSES' DIRECTORY

DISTRICT NO. 1

ARIZONA STATE NURSES ASS'N

**MRS. MARJORIE E. KASUN, R.N.,
Registrar**

Nurses' Professional Registry

10 S. 12th Ave. Phoenix Ph. ALpine 4-4151

MARICOPA COUNTY MEDICAL SOCIETY DIRECTORY

Operated by District No. 1 of the
Arizona State Nurses' Association

Mrs. Marjorie E. Kasun, R.N., Director
AL 3-5101

Office Open 24 Hours Daily
10 South 12th Avenue — Phoenix

PHYSICIANS' DIRECTORY

NEUROLOGY and PSYCHIATRY

OTTO L. BENDHEIM, M.D.

NEUROLOGY and PSYCHIATRY

1515 North Ninth Street
PHOENIX, ARIZONA

Certified by American Board of
Psychiatry and Neurology

ROBERT L. BEAL, M.D.

Practice Limited to Psychiatry and Neurology

Park Central Medical Building
550 W. Thomas Road — 234 Patio D
Phone CR 4-6711
Phoenix, Arizona

RICHARD E. H. DUISBERG, M.D.

Diplomate American Board of Psychiatry and
Neurology

Phoenix, Arizona

THIS SPACE FOR SALE
FOR INFORMATION AND RATES
write to

ARIZONA MEDICINE

424 Heard Bldg.
Phone ALpine 2-4884
PHOENIX, ARIZONA

PROCTOLOGY

WALLACE M. MEYER, M.D.

PROCTOLOGY

Park Central Medical Bldg.

Phone CR 4-5632
550 W. Thomas Road — 216 Patio B
Phoenix, Arizona

Plastic and Reconstructive Surgery

HOWARD C. LAWRENCE, M.D.

Diplomate of the
American Board of Plastic Surgery

709 Professional Building
15 E. Monroe Street Phone ALpine 8-4101
Phoenix, Arizona

PHYSICIANS' DIRECTORY

UROLOGY**ROBERT H. CUMMINGS, M.D.**

Diplomate of the American
Board of Urology

Park Central Medical Bldg.
Phone CR 4-4912
550 W. Thomas Road — 207 Patio A
Phoenix, Arizona

W. G. SHULTZ, M.D., F. A. C. S.

Diplomate of The American
Board of Urology

E. R. UPDEGRAFF, M.D.

1010 N. Country Club Road
Telephone 5-2609 Tucson, Arizona

PAUL L. SINGER, M.D., F. A. C. S.

Certified American Board of
UROLOGY

1313 N. Second Street Phone ALpine 3-1739
PHOENIX, ARIZONA

DONALD B. LEWIS, M.D.

UROLOGY

Certified by the American Board of Urology
123 So. Stone Ave. Phone 2-7081
Tucson, Arizona

ALLERGY**H. M. PURCELL, M.D.**

Diplomate of the American Board of Urology
Park Central Medical Building
550 W. Thomas Road
Phone CR 4-5202
Phoenix, Arizona

E. A. GATTERDAM, M.D.

ALLERGY

15 E. Monroe St., Professional Bldg.
Office Hours: 11 A.M. to 5 P.M.
Phoenix, Arizona

CLINIC**NELSON CLINIC**

D. E. NELSON, M.D.
A. H. ERICKSON, MD.
D. CLUFF, D.D.S.

503 Fifth Avenue
SAFFORD, ARIZONA

MALIGNANT DISEASE**JAMES M. OVENS, M.D.**

F.A.C.S. F.I.C.S.

Diplomate American Board of Surgery
Cancer and Allied Diseases
608 Professional Bldg. Phone ALpine 4-1973
Phoenix, Arizona

ANESTHESIOLOGY

LOUISE BEWERSDORF, M.D.
HERMAN BEWERSDORF, M.D.

ANESTHESIOLOGY
Park Central Medical Bldg.
Phone CR 4-5674
550 W. Thomas Road — 24 R
Phoenix, Arizona

HOSPITAL**H. B. LEHMBERG, M.D.****J. T. O'NEIL, M.D.**

Casa Grande Clinic * Phone 4495
Casa Grande, Arizona

PHYSICIANS' DIRECTORY

INTERNAL MEDICINE**ROBERT S. FLINN, M.D.**INTERNAL MEDICINE
CARDIOLOGY and ELECTROCARDIOGRAPHY

Park Central Medical Bldg.

Phone AM 6-8485

550 W. Thomas Road — 217 Patio B

Phoenix, Arizona

JESSE D. HAMER, M.D.**F. A. C. P.**

INTERNAL MEDICINE

CARDIOLOGY

Suit 910
15 E. Monroe St.Phoenix
Arizona**FRANK J. MILLOY, M.D.**
F. A. C. P.Diplomate of the American Board of
Internal Medicine

INTERNAL MEDICINE

611 Professional Building

Phone ALpine 4-2171

Phoenix, Arizona

DAVID M. MARCUS, M.D.

INTERNAL MEDICINE

1850 N. Laurel Avenue — Phone ALpine 4-7970

Phoenix, Arizona

ROBERT E. RIDER, M.D.INTERNAL MEDICINE
ELECTROCARDIOGRAPHY

Del Sol Hotel Bldg.

Phone 3-3721

Yuma, Arizona

JOSEPH BANK, M.D.Diplomate of
American Board of Internal Medicine
American Board of Gastroenterology

GASTROENTEROLOGY, GASTROSCOPY

800 North First Avenue Phone: ALpine 4-7245

PHOENIX, ARIZONA

OBSTETRICS and GYNECOLOGY**HARRY J. FELCH, M.D.**

Physician and Surgeon

Residence

325 W. Granada
Phoenix, Arizona
Res. AL 3-1151

Office

703 Professional Bldg.
15 E. Monroe Street
Office AL 3-1151**GYNECOLOGY & ENDOCRINOLOGY****JOSEPH B. RADDIN, M. D.**Practice limited to
MEDICAL GYNECOLOGY & ENDOCRINOLOGY

619 Professional Building

15 E. Monroe — Phoenix, Arizona

Phone ALpine 2-3577

CHILDREN'S DISEASES**WM. F. SCHOFFMAN, M.D.**
CECELIA H. SHEMBAB, M.D.
JOHN R. KEEFREY, M.D.

DOCTORS BUILDING

316 W. McDowell Rd. — Telephone ALpine 4-7287
Phoenix, Arizona**DERMATOLOGY****GEORGE K. ROGERS, M.D.**

DERMATOLOGY

Diplomate of American Board of
Dermatology and Syphilology

Phone ALpine 3-5264

105 W. McDowell Road Phoenix, Arizona

PHYSICIANS' DIRECTORY

SURGERY

**EDWARD L. KETTENBACH, M.D.,
F.A.C.S., F.I.C.S.**
SURGERY

Diplomate American Board of Surgery
2324 North Tucson Blvd. Phone 5-2605
Tucson, Arizona

**DELBERT L. SECRIST, M.D.,
F. A. C. S.**

123 South Stone Avenue
Tucson, Arizona
Office Phone 2-3371 Home Phone 5-9433

H. D. KETCHERSIDE, M.D.
SURGERY and UROLOGY

800 North First Avenue
Phone ALpine 4-7245
Phoenix, Arizona

W. R. MANNING, M.D., F.A.C.S.
SURGERY

Diplomate American Board of Surgery
620 North Country Club Road Phone 5-2687
Tucson, Arizona

DONALD A. POLSON, M.D., M. Sc.
GENERAL SURGERY

Certified by the American Board of Surgery
550 W. Thomas Road
Phone CRestwood 4-2081
Phoenix, Arizona

**THOMAS H. BATE, M.D.
F.A.C.S., F.I.C.S.M.Sc. (Surgery)**
PRACTICE LIMITED TO SURGERY

Diplomate American Board of Surgery
15 E. Monroe — Office Phone ALpine 4-3326
Phoenix, Arizona

ORTHOPEDIC SURGERY

**GEORGE L. DIXON, M.D.
PHILIP G. DERICKSON, M.D.**
ORTHOPAEDIC SURGERY

Diplomates of the American Board
of Orthopaedic Surgery
744 N. Country Club Road Telephone 5-1533
TUCSON, ARIZONA

**GEO. A. WILLIAMSON, M.D.,
F. A. C. S.**

Diplomate American Board of Orthopaedic Surgery

LEO L. TUVESON, M.D.

Orthopaedic Surgery
Park Central Medical Building
550 West Thomas Road — 116 Patio C
Telephone CRestwood 4-5459 — Phoenix, Arizona

**ROBERT E. HASTINGS, M.D.,
F. A. C. S.**

Diplomate American Board of Orthopaedic
Surgery

ROBERT W. WEBER, M.D.

ORTHOPAEDIC SURGERY
1014 N. Country Club
TUCSON, ARIZONA

**STANLEY S. TANZ, M.D., F.A.C.S.
ORTHOPAEDIC SURGERY**

Diplomate American Board of Orthopaedic Surgery
2530 East Broadway Telephone 5-0114
Tucson, Arizona

